arXiv:1210.1846v1 [math.NA] 6 Oct 2012

Convergence Rate and Quasi-Optimal Complexity of Adaptive Finite Element Computations for Multiple Eigenvalues *

Xiaoying Dai† Lianhua He‡ Aihui Zhou§

Abstract

In this paper, we study an adaptive finite element method for multiple eigenvalue problems. We obtain both convergence rate and quasi-optimal complexity of the adaptive finite element eigenvalue approximation, without any additional assumption to those required in the adaptive finite element analysis for the boundary value problem. Our analysis is based on a certain relationship between the finite element eigenvalue approximation and the associated finite element boundary value approximation and a crucial property of eigenspace approximation which are also presented in the paper.

Key words. Adaptive finite element, a posteriori error estimator, complexity, convergence, multiple eigenvalue.
2000 AMS subject classifications. 65F15, 65N15, 65N25, 65N30, 65N50

1 Introduction

Adaptive finite element computation is one of efficient approaches to solve partial differential equations and has been successfully used in scientific and engineering computing. Its numerical analysis has been also derived much attention from the mathematical community. Since Babuška and Vogelius [4] gave an analysis of an adaptive finite element method (AFEM) for linear symmetric elliptic problems in one dimension, there has been much investigation on the convergence and complexity of AFEMs in literature (see, e.g., [6, 7, 12, 13, 16, 31, 32] and the references cited therein). In the context of the finite element approximations of eigenvalue problems, in particular, we note that there are a number of works concerning a posteriori error estimates [5, 14, 21, 22, 24, 33], AFEM convergence [12, 15, 16, 17, 20] and complexity [12, 15, 20]. Except for the convergence analysis in [16], to our best knowledge, there is no any work about convergence rate and complexity of AFEM for multiple eigenvalue problems. The purpose of this paper is to fill in the gap.

We understand that multiple eigenvalue problems are topic in science and engineering, such as Hartree-Fock equation and Kohn-Sham equation used to model ground state electronic structures of

---

[*] This work was partially supported by the Funds for Creative Research Groups of China under Grant 11021101, the National Basic Research Program of China under Grant 2011CB309703, the National Science Foundation of China under Grant 11101416, and the National Center for Mathematics and Interdisciplinary Sciences, Chinese Academy of Sciences.

[†] LSEC, Institute of Computational Mathematics and Scientific/Engineering Computing, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China (daixy@lsec.cc.ac.cn).

[‡] LSEC, Institute of Computational Mathematics and Scientific/Engineering Computing, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China (helh@lsec.cc.ac.cn).

[§] LSEC, Institute of Computational Mathematics and Scientific/Engineering Computing, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China (azhou@lsec.cc.ac.cn).

molecular systems in quantum chemistry and materials science, in which hundreds of thousands of eigenvalues and their corresponding eigenfunctions are needed, and among these eigenvalues, most are multiple [23, 25, 29, 30, 11]. While the central computation in solving Hartree-Fock equation or Kohn-Sham equation is the repeated solution of linear Schrödinger type equation, of which adaptive finite element analysis and computation are significant. Hence, we want to study the convergence and complexity of AFEMs for multiple eigenvalue problems and focus on the following elliptic eigenvalue problems: find $\lambda \in \mathbb{R}$ and $u \in H_0^1(\Omega)$ such that

$$\left\{ \begin{array}{rll} -\nabla \cdot (A\nabla u) + cu & = \lambda u & \text{in} \quad \Omega, \\ \|u\|_{0,\Omega} & = 1 & \text{on} \quad \partial\Omega, \end{array} \right.$$

where A, c are coefficients stated precisely in Section 2.

We see that the analysis technologies for the convergence rate and complicity of AFEM in literature are only valid for simple eigenvalues and their corresponding eigenfunctions, it can not be applied directly to multiple eigenvalue cases. The difficulty lies in that in context of multiple eigenvalue cases, it is not practicable to figure out the discreted eigenfunctions obtained over different meshes so as to approximate the same exact eigenfunctions. As a result, the standard technology of measuring the error of every eigenfunction does not work well any more, which results in the difficulty when analyzing the reduction for error of the approximate eigenfunction over two consecutive meshes. Instead, we employ some measure between the eigenspace, the space of eigenfunctions, and its approximation. By using the similar perturbation argument in [12, 20] together with eigenspace approximation technology and its crucial property that is also shown in this paper (see Lemma 3.1), we obtain the convergence rate and quasi-optimal complexity of AFEM for multiple eigenvalue problems.

The paper is organized as follows. In the next section, we shall describe some basic notation and review the existing results of finite element approximations to a class of linear second order elliptic boundary value and eigenvalue problems, which will be used in our analysis. In Section 3, we construct the a posteriori error estimators for finite element eigenvalue problems from the relationship between the elliptic eigenvalue approximation and the associated boundary value approximation and then design adaptive finite element algorithm for the elliptic eigenvalue problems. We analyze the convergence and quasi-optimal complexity of the adaptive finite element eigenvalue computations in Sections 4 and 5, respectively. We present several numerical examples in Section 6 to illustrate our theory. Finally, we remark how our main results can be expected for the Steklov eigenvalue problems and inexact numerical solutions.

2 Preliminaries

Let $\Omega \subset \mathbb{R}^d (d \geq 1)$ be a polytopic bounded domain. We shall use the standard notation for Sobolev spaces $W^{s,p}(\Omega)$ and their associated norms and seminorms, see, e.g., [1, 9]. For $p = 2$, we denote $H^s(\Omega) = W^{s,2}(\Omega)$ and $H_0^1(\Omega) = \{v \in H^1(\Omega) : v \mid_{\partial\Omega} = 0\}$, where $v \mid_{\partial\Omega} = 0$ is understood in the sense of trace, $\| \cdot \|_{s,\Omega} = \| \cdot \|_{s,2,\Omega}$, and $(\cdot, \cdot)$ is the standard L^2 inner product. Throughout this paper, we shall use C to denote a generic positive constant which may stand for different values at its different occurrences. For convenience, the symbol $\lesssim$ will be used in this paper. The notation that $A \lesssim B$ means that $A \leq CB$ for some constant C that is independent of mesh parameters. All the constants involved are independent of mesh sizes.

Let $\{\mathcal{T}_h\}$ be a shape regular family of nested conforming meshes over Ω: there exists a constant γ^* such that

$$\frac{h_T}{\rho_T} \leq \gamma^* \quad \forall T \in \bigcup_h \mathcal{T}_h,$$

where, for each $T \in \mathcal{T}_h$, h_T is the diameter of T, and ρ_T is the diameter of the biggest ball contained in $T, h = \max\{h_T : T \in \mathcal{T}_h\}$. Let $\mathcal{E}_h$ denote the set of interior faces (edges or sides) of $\mathcal{T}_h$.

Let $S^{h,k}(\Omega)$ be a space of continuous functions on Ω such that for $v \in S^{h,k}(\Omega)$, v restricted to each T is a polynomial of degree not greater than k, namely,

$$S^{h,k}(\Omega) = \{v \in C(\bar{\Omega}) : \ v|_T \in P_T^k \quad \forall T \in \mathcal{T}_h\},$$

where P_T^k is the space of polynomials of degree not greater than a positive integer k. Set $S_0^{h,k}(\Omega) = S^{h,k}(\Omega) \cap H_0^1(\Omega)$. We shall denote $S_0^{h,k}(\Omega)$ by $S_0^h(\Omega)$ for simplification of notation afterwards.

2.1 A linear elliptic boundary value problem

In this subsection, we shall present some basic properties of a second order elliptic boundary value problem and its finite element approximations.

Consider the homogeneous boundary value problem:

$$\begin{cases} Lu_i &= f_i \ \text{in} \quad \Omega, \quad i = 1, \cdots, N, \\ \ \ u_i &= 0 \ \text{on} \quad \partial\Omega, \end{cases} \tag{2.1}$$

where N is a positive integer, L is a linear second order elliptic operator:

$$Lu = -\nabla \cdot (A\nabla u) + cu$$

with $A : \Omega \to \mathbb{R}^{d\times d}$ being piecewise Lipschitz over initial triangulation and symmetric positive definite with smallest eigenvalue uniformly bounded away from 0, and $0 \le c \in L^\infty(\Omega)$.

The weak form of (2.1) reads: find $U \equiv (u_1, \cdots, u_N) \in (H_0^1(\Omega))^N$ such that

$$a(u_i, v_i) = b(f_i, v_i) \qquad \forall v_i \in H_0^1(\Omega), i = 1, \cdots, N, \tag{2.2}$$

where

$$a(u, v) = (A\nabla u, \nabla v) + (cu, v) \quad \text{and} \quad b(u, v) = (u, v).$$

We observe that $a(\cdot, \cdot)$ is a bounded bilinear form over $H_0^1(\Omega)$:

$$|a(w, v)| \le C_a \|w\|_{1,\Omega} \|v\|_{1,\Omega} \quad \forall w, v \in H_0^1(\Omega)$$

and for energy norm $\|\cdot\|_{a,\Omega}$, which is defined by $\|w\|_{a,\Omega} = \sqrt{a(w, w)}$, there hold

$$c_a \|w\|_{1,\Omega} \le \|w\|_{a,\Omega} \le C_a \|w\|_{1,\Omega} \quad \forall w \in H_0^1(\Omega),$$

where c_a and C_a are positive constants. We understand that (2.2) is uniquely solvable for any $f_i \in H^{-1}(\Omega) (i = 1, \cdots, N)$.

For $L^2(\Omega)$ with $\|\cdot\|_{b,\Omega} = \sqrt{b(\cdot, \cdot)}$, we see that there is a unique compact operator $K : L^2(\Omega) \to H_0^1(\Omega)$ satisfying

$$a(Kw, v) = b(w, v) \quad \forall w \in L^2(\Omega), v \in H_0^1(\Omega).$$

Define the Galerkin-projection $R_h : H_0^1(\Omega) \to V_h \equiv S_0^h(\Omega)$ by

$$a(u - R_h u, v) = 0 \quad \forall u \in H_0^1(\Omega) \ \forall v \in V_h, \tag{2.3}$$

then apparently

$$\|R_h u\|_{1,\Omega} \lesssim \|u\|_{1,\Omega} \quad \forall u \in H_0^1(\Omega).$$

The following results can be found in [2, 34].

Proposition 2.1. *Let*

$$\rho_{\Omega}(h) = \sup_{f\in L^2(\Omega),\|f\|_{b,\Omega}=1} \inf_{v\in V_h} \|Kf - v\|_{1,\Omega}.$$

Then $\rho_{\Omega}(h) \to 0$ *as* $h \to 0$ *and*

$$\|u - R_h u\|_{b,\Omega} \lesssim \rho_{\Omega}(h)\|u - R_h u\|_{1,\Omega} \quad \forall u \in H_0^1(\Omega).$$

A standard finite element scheme for (2.2) reads: find $U_h = (u_{1,h}, \cdots, u_{N,h}) \in V_h^N$ satisfying

$$a(u_{i,h}, v_i) = b(f_i, v_i) \ \ \forall v_i \in V_h, i = 1, \cdots, N. \tag{2.4}$$

Let $\mathbb{T}$ denote the class of all conforming refinements by bisection of $\mathcal{T}_{h_0}$. For $\mathcal{T}_h \in \mathbb{T}$, define the element residual $\tilde{\mathcal{R}}_T(u_{i,h})$ and the jump residual $\tilde{J}_E(u_{i,h})$ by

$$\begin{aligned}\tilde{\mathcal{R}}_T(u_{i,h}) &= f_i - Lu_{i,h} = f_i + \nabla\cdot(A\nabla u_{i,h}) - cu_{i,h} \quad \text{in } T \in \mathcal{T}_h,\\ \tilde{J}_E(u_{i,h}) &= -A\nabla u_{i,h}^+ \cdot \nu^+ - A\nabla u_{i,h}^- \cdot \nu^- = [[A\nabla u_{i,h}]]_E \cdot \nu_E \quad \text{on } E \in \mathcal{E}_h,\end{aligned}$$

where E is the common side of elements T^+ and T^- with unit outward normals ν^+ and ν^-, respectively, and $\nu_E = \nu^-$. Let ω_T be the union of elements sharing a side with T and ω_E be the union of elements which shares the side E, that is, $\omega_E = T^+ \cap T^-$.

For $T \in \mathcal{T}_h$, we define the local error indicator $\tilde{\eta}_h(u_{i,h}, T)$ by

$$\tilde{\eta}_h^2(u_{i,h}, T) = h_T^2\|\tilde{\mathcal{R}}_T(u_{i,h})\|_{0,T}^2 + \sum_{E\in\mathcal{E}_h, E\subset\partial T} h_E\|\tilde{J}_E(u_{i,h})\|_{0,E}^2$$

and the oscillation $\widetilde{osc}_h(u_{i,h}, T)$ by

$$\widetilde{osc}_h^2(u_{i,h}, T) = h_T^2\|\tilde{\mathcal{R}}_T(u_{i,h}) - \overline{\tilde{\mathcal{R}}_T(u_{i,h})}\|_{0,T}^2 + \sum_{E\in\mathcal{E}_h, E\subset\partial T} h_E\|\tilde{J}_E(u_{i,h}) - \overline{\tilde{J}_E(u_{i,h})}\|_{0,E}^2,$$

where $\overline{w}$ is the L^2-projection of $w \in L^2(\Omega)$ to polynomials of some degree on T or E.

We define the error estimator $\tilde{\eta}_h(u_{i,h}, \Omega)$ and the oscillation $\widetilde{osc}_h(u_{i,h}, \Omega)$ by

$$\tilde{\eta}_h^2(u_{i,h}, \Omega) = \sum_{T\in\mathcal{T}_h, T\subset\Omega} \tilde{\eta}_h^2(u_{i,h}, T) \quad \text{and} \quad \widetilde{osc}_h^2(u_{i,h}, \Omega) = \sum_{T\in\mathcal{T}_h, T\subset\Omega} \widetilde{osc}_h^2(u_{i,h}, T).$$

For any $U_h = (u_{1,h}, \cdots, u_{N,h}) \in V_h^N$ we set

$$\tilde{\eta}_h^2(U_h, \Omega) = \sum_{i=1}^N \tilde{\eta}_h^2(u_{i,h}, \Omega) \quad \text{and} \quad \widetilde{osc}_h^2(U_h, \Omega) = \sum_{i=1}^N \widetilde{osc}_h^2(u_{i,h}, \Omega).$$

In our analysis we need the following lemma [7].

Lemma 2.1. *There exists a constant* C_* *which depends on* A*, regularity constant* γ^**, and coefficient* c*, such that*

$$\widetilde{osc}_h(v, T) \le \widetilde{osc}_h(w, T) + C_*\|v - w\|_{1,\omega_T}, \quad \forall v, w \in V_h. \tag{2.5}$$

We have the standard a posteriori error estimates for the finite element approximation of boundary value problems (2.1) as follows (c.f., e.g., [27, 28, 33])

$$\|u_i - u_{i,h}\|_{a,\Omega} \le \tilde{C}_1 \tilde{\eta}_h(u_{i,h}, \Omega), \tag{2.6}$$

$$\tilde{C}_2^2 \tilde{\eta}_h^2(u_{i,h}, \Omega) - \tilde{C}_3^2 \widetilde{osc}_h^2(u_{i,h}, \Omega) \le \|u_i - u_{i,h}\|_{a,\Omega}^2, \tag{2.7}$$

where $\tilde{C}_1, \tilde{C}_2$ and $\tilde{C}_3$ are positive constants depending on the shape regularity of the mesh $\mathcal{T}_h$.

The adaptive algorithm for solving (2.4) without oscillation marking can be stated as follows.

Algorithm C_0

Choose a parameter $0 < \theta < 1$.

1. Pick any initial mesh $\mathcal{T}_{h_0}$.
2. Solve the system on $\mathcal{T}_{h_0}$ for the discrete solution $U_{h_0} \equiv (u_{1,h_0}, \cdots, u_{N,h_0})$.
3. Let $k = 0$.
4. Compute the local indicators $\tilde{\eta}_{h_k}$.
5. Construct $\mathcal{M}_{h_k} \subset \mathcal{T}_{h_k}$ by **Marking Strategy** E_0 and parameter θ.
6. Refine $\mathcal{T}_{h_k}$ to get a new conforming mesh $\mathcal{T}_{h_{k+1}}$ by Procedure **REFINE**.
7. Solve the system on $\mathcal{T}_{h_{k+1}}$ to get the discrete solution $U_{h_{k+1}} \equiv (u_{1,h_{k+1}}, \cdots, u_{N,h_{k+1}})$.
8. Let $k = k + 1$ and go to Step 4.

Marking Strategy E_0 in **Algorithm** C_0 was introduced [13, 28] when $N = 1$. It is used to enforce error reduction and can be defined as follows.

Marking Strategy E_0

Given a parameter $0 < \theta < 1$.

1. Construct a minimal subset $\mathcal{M}_{h_k}$ of $\mathcal{T}_{h_k}$ by selecting some elements in $\mathcal{T}_{h_k}$ such that

$$\sum_{T \in \mathcal{M}_{h_k}} \tilde{\eta}_{h_k}^2(U_{h_k}, T) \ge \theta \tilde{\eta}_{h_k}^2(U_{h_k}, \Omega).$$

2. Mark all the elements in $\mathcal{M}_{h_k}$.

The procedure **REFINE** here is not required to satisfy the Interior Node Property of [27, 28]. Given a fixed number $b \geq 1$, for any $\mathcal{T}_{h_k} \in \mathbb{T}$ and a subset $\mathcal{M}_{h_k} \subset \mathcal{T}_{h_k}$ of marked elements,

$$\mathcal{T}_{h_{k+1}} = \textbf{REFINE}(\mathcal{T}_{h_k}, \mathcal{M}_{h_k})$$

outputs a conforming triangulation $\mathcal{T}_{h_{k+1}} \in \mathbb{T}$, where at least all elements of $\mathcal{M}_{h_k}$ are bisected b times. We define $R_{\mathcal{T}_{h_k} \to \mathcal{T}_{h_{k+1}}} = \mathcal{T}_{h_k} \backslash (\mathcal{T}_{h_k} \cap \mathcal{T}_{h_{k+1}})$ as the set of refined elements, thus $\mathcal{M}_{h_k} \subset R_{\mathcal{T}_{h_k} \to \mathcal{T}_{h_{k+1}}}$.

Lemma 2.2. *Assume that $\mathcal{T}_{h_0}$ verifies condition (b) of Section 4 in [32]. Let $\{\mathcal{T}_{h_k}\}_{k\geq 0}$ be any sequence of refinements of $\mathcal{T}_{h_0}$ where $\mathcal{T}_{h_{k+1}}$ is generated from $\mathcal{T}_{h_k}$ by $\mathcal{T}_{h_{k+1}} = \textbf{REFINE}(\mathcal{T}_{h_k}, \mathcal{M}_{h_k})$ with a subset $\mathcal{M}_{h_k} \subset \mathcal{T}_{h_k}$. Then*

$$\#\mathcal{T}_{h_k} - \#\mathcal{T}_{h_0} \lesssim \sum_{j=0}^{k-1} \#\mathcal{M}_{h_j} \quad \forall k \geq 1$$

is valid, where the hidden constant depends on $\mathcal{T}_{h_0}$ and b.

By some primary operation, we can easily extend the results about the case of $N = 1$ in [7] to vector problems, and obtain the following results.

Theorem 2.1. *Let $\{U_{h_k}\}_{k\in\mathbb{N}_0}$ be a sequence of finite element solutions of boundary problems produced by* **Algorithm** C_0*. Then there exist constants $\tilde{\gamma} > 0$ and $\xi \in (0, 1)$, depending only on the shape regularity of meshes, the data, and the parameters used by* **Algorithm** C_0*, such that for any two consecutive iterates k and $k + 1$ we have*

$$\|U - U_{h_{k+1}}\|^2_{a,\Omega} + \tilde{\gamma}\tilde{\eta}^2_{h_{k+1}}(U_{h_{k+1}}, \Omega) \leq \xi^2 \Big(\|U - U_{h_k}\|^2_{a,\Omega} + \tilde{\gamma}\tilde{\eta}^2_{h_k}(U_{h_k}, \Omega)\Big).$$

Indeed, the constant $\tilde{\gamma}$ has the following form

$$\tilde{\gamma} = \frac{1}{(1 + \delta^{-1})C_*^2} \tag{2.8}$$

with some constant $\delta \in (0, 1)$.

Lemma 2.3. *Let $u_{h_k,l} \in V_{h_k}$ and $u_{h_{k+1},l} \in V_{h_{k+1}}(l = 1, \cdots, N)$ be discrete solutions of (2.4) over a conforming mesh $\mathcal{T}_{h_k}$ and its refinement $\mathcal{T}_{h_{k+1}}$ with marked element $\mathcal{M}_{h_k}$. Let $\mathcal{R} = R_{\mathcal{T}_{h_k} \to \mathcal{T}_{h_{k+1}}}$ be the set of refined elements, then the following localized upper bound is valid*

$$\|U_{h_k} - U_{h_{k+1}}\|^2_{a,\Omega} \leq \tilde{C}_1^2 \sum_{T\in\mathcal{R}} \tilde{\eta}^2_{h_k}(U_{h_k}, T),$$

where $U_{h_k} \equiv (u_{h_k,1} \cdots, u_{h_k,q})$ and $U_{h_{k+1}} \equiv (u_{h_{k+1},1} \cdots, u_{h_{k+1},q})$.

Proposition 2.2. *Let $u_{h_k,l} \in V_{h_k}$ and $u_{h_{k+1},l} \in V_{h_{k+1}}(l = 1, \cdots, N)$ be discrete solutions of (2.4) over a conforming mesh $\mathcal{T}_{h_k}$ and its refinement $\mathcal{T}_{h_{k+1}}$ with marked element $\mathcal{M}_{h_k}$. Suppose that they satisfy the energy decrease property*

$$\|U - U_{h_{k+1}}\|^2_{a,\Omega} + \tilde{\gamma}_0\widetilde{osc}^2_{h_{k+1}}(U_{h_{k+1}}, \Omega) \leq \tilde{\xi}_0^2\big(\|U - U_{h_k}\|^2_{a,\Omega} + \tilde{\gamma}_0\widetilde{osc}^2_{h_k}(U_{h_k}, \Omega)\big)$$

with $\tilde{\gamma}_0 > 0$ being a constant and $\tilde{\xi}_0^2 \in (0, \frac{1}{2})$. Then the set $\mathcal{R} = R_{\mathcal{T}_{h_k} \to \mathcal{T}_{h_{k+1}}}$ satisfies the Dörfler property

$$\sum_{T\in\mathcal{R}} \tilde{\eta}^2_{h_k}(U_{h_k}, T) \geq \tilde{\theta} \sum_{T\in\mathcal{T}_{h_k}} \tilde{\eta}^2_{h_k}(U_{h_k}, T)$$

with $\tilde{\theta} = \frac{\tilde{C}_2^2(1-2\tilde{\xi}_0^2)}{\tilde{C}_0(\tilde{C}_1^2+(1+2C_*^2\tilde{C}_1^2)\tilde{\gamma}_0)}$, where $\tilde{C}_0 = \max(1, \frac{\tilde{C}_3^2}{\tilde{\gamma}_0})$.

2.2 A linear eigenvalue problem

A number λ is called an eigenvalue of the form $a(\cdot,\cdot)$ relative to the form $b(\cdot,\cdot)$ if there is a nonzero function $0 \neq u \in H_0^1(\Omega)$, called an associated eigenfunction, satisfying

$$a(u,v) = \lambda b(u,v) \quad \forall v \in H_0^1(\Omega). \tag{2.9}$$

We see that (2.9) has a countable sequence of real eigenvalues

$$0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \cdots$$

and corresponding eigenfunctions

$$u_1, u_2, u_3, \cdots,$$

which can be assumed to satisfy

$$b(u_i, u_j) = \delta_{ij}, \ i,j = 1,2,\cdots$$

In the sequence $\{\lambda_j\}$, the λ_j's are repeated according to geometric multiplicity.

The following property of eigenvalue and eigenfunction approximation is useful (see [3, 35]).

Proposition 2.3. *Let (λ, u) be an eigenpair of (2.9). For any $w \in H_0^1(\Omega) \setminus \{0\}$,*

$$\frac{a(w,w)}{b(w,w)} - \lambda = \frac{a(w-u,w-u)}{b(w,w)} - \lambda \frac{b(w-u,w-u)}{b(w,w)}.$$

A standard finite element scheme for (2.9) is: find a pair of (λ_h, u_h), where λ_h is a number and $0 \neq u_h \in V_h$, satisfying

$$a(u_h, v) = \lambda_h b(u_h, v) \quad \forall v \in V_h. \tag{2.10}$$

Let us order the eigenvalues of (2.10) as follows

$$0 < \lambda_{1,h} \leq \lambda_{2,h} \leq \cdots \leq \lambda_{n_h,h}, \ \ n_h = \dim V_h,$$

and assume the corresponding eigenfunctions

$$u_{1,h}, u_{2,h}, \cdots, u_{n_h,h}$$

satisfy

$$b(u_{i,h}, u_{j,h}) = \delta_{ij}, \ i,j = 1,2,\cdots,n_h.$$

As a consequence of the minimum-maximum principle (see [3] or [8]) and Proposition 2.3, we have

$$\lambda_i \leq \lambda_{i,h} \leq \lambda_i + C_i \|u_i - u_{i,h}\|_a^2, \ \ i = 1,2,\cdots,n_h. \tag{2.11}$$

Let λ be any eigenvalue of (2.9) with multiplicity q and $M(\lambda)$ denote the space of eigenfunctions corresponding to λ, that is

$$M(\lambda) = \{w \in H_0^1(\Omega) : w \text{ is an eigenvector of (2.9) corresponding to } \lambda\}.$$

We see that λ will be approximated from above by the Galerkin approximate eigenvalues:

$$\lambda \leq \lambda_{h,1} \leq \cdots \leq \lambda_{h,q}.$$

Set

$$\delta_h(\lambda) = \sup_{w \in M(\lambda), \|w\|_{b,\Omega}=1} \inf_{v \in V_h} \|w - v\|_{a,\Omega}.$$

The following results are standard and can be found in literature (see, e.g., [2, 3, 8] or [35]).

Proposition 2.4. *Let λ be any eigenvalue of (2.9) with multiplicity q and $u_{h,1},\cdots,u_{h,q}$ with $\|u_{h,l}\|_{b,\Omega}=1 (l=1,\cdots,q)$ be the Galerkin eigenfunctions corresponding to $\lambda_{h,1},\cdots,\lambda_{h,q}$, respectively. There hold*

$$\|u-E_hu\|_{b,\Omega}\lesssim\rho_\Omega(h)\|u-E_hu\|_{a,\Omega}\quad\forall u\in M(\lambda),\tag{2.12}$$

$$\|u_{h,l}-Eu_{h,l}\|_{b,\Omega}\lesssim\rho_\Omega(h)\|u_{h,l}-Eu_{h,l}\|_{a,\Omega},\tag{2.13}$$

where E denotes the orthogonal projection of the energy space onto $M(\lambda)$, E_h the orthogonal projection onto span$\{u_{h,1},\cdots,u_{h,q}\}$.

For convenience, here and hereafter, we let $M_h(\lambda)=\mathrm{span}\{u_{h,1},\cdots,u_{h,q}\}$.

3 Adaptive finite element method

Here and hereafter we assume that λ is some eigenvalue of (2.9) with multiplicity q and $u\in M(\lambda)$, and $(\lambda_{h,l},u_{h,l})\in\mathbb{R}\times V_h$ is the associated finite element eigenpairs of (2.10) with $\|u_{h,l}\|_{b,\Omega}=1(l=1,\cdots,q)$ that satisfy (2.12) and (2.13).

Let $\lambda^h=a(E_hu,E_hu)$, we conclude from (2.11) and (2.12) that

$$\|u-E_hu\|_{b,\Omega}+|\lambda-\lambda^h|\le C(\rho_\Omega(h)+\|u-E_hu\|_{a,\Omega})\|u-E_hu\|_{a,\Omega}.\tag{3.1}$$

Note that (2.9) and (2.10) can be rewritten as

$$u=\lambda Ku,\quad u_h=\lambda_hR_hKu_h,$$

where R_h is the Galerkin projection defined by (2.3).

We see from the definition of E_h that for any $u\in M(\lambda)$, there exist some constants $\{\alpha_{h,l}(u)\}_{l=1}^q$ satisfying $\sum_{l=1}^q\big(\alpha_{h,l}(u)\big)^2=1$ such that $E_hu=\sum_{l=1}^q\alpha_{h,l}(u)u_{h,l}$. Define $w^h=\sum_{l=1}^q\alpha_{h,l}(u)\lambda_{h,l}Ku_{h,l}$, and we obtain from a direct calculation that

$$E_hu=R_hw^h.\tag{3.2}$$

Theorem 3.1. *Given $u\in M(\lambda)$, let $r(h)=\rho_\Omega(h)+\|u-E_hu\|_{a,\Omega}$. Then*

$$\|u-E_hu\|_{a,\Omega}=\|w^h-R_hw^h\|_{a,\Omega}+\mathcal{O}(r(h))\|u-E_hu\|_{a,\Omega}.\tag{3.3}$$

Proof. We obtain from the definition of w^h that

$$\begin{aligned}u-w^h&=\lambda Ku-\sum_{l=1}^q\alpha_{h,l}(u)\lambda_{h,l}Ku_{h,l}\\&=\lambda K(u-E_hu)+\lambda K(\sum_{l=1}^q\alpha_{h,l}(u)u_{h,l})-\sum_{l=1}^q\alpha_{h,l}(u)\lambda_{h,l}Ku_{h,l}\\&=\lambda K(u-E_hu)+\sum_{l=1}^q\alpha_{h,l}(u)(\lambda-\lambda_{h,l})Ku_{h,l}.\end{aligned}$$

Since

$$\lambda-\lambda^h=\sum_{l=1}^q\big(\alpha_{h,l}(u)\big)^2(\lambda-\lambda_{h,l}),$$

we have

$$\begin{aligned}\|\sum_{l=1}^{q}\alpha_{h,l}(u)(\lambda-\lambda_{h,l})Ku_{h,l}\|_{a,\Omega} &\lesssim |\sum_{l=1}^{q}\alpha_{h,l}(u)(\lambda-\lambda_{h,l})|\\ &\leq \max_{\alpha_{h,l}(u)\neq 0,l=1,\cdots,q}\{\frac{1}{|\alpha_{h,l}(u)|}\}|\sum_{l=1}^{q}\alpha_{h,l}^2(\lambda-\lambda_{h,l})|\\ &\lesssim |\lambda-\lambda^h|,\end{aligned}$$

which together with the fact $\|K(u-E_hu)\|_{a,\Omega}\lesssim\|u-E_hu\|_{b,\Omega}$ and (3.1) leads to

$$\|u-w^h\|_{a,\Omega} \quad\leq\quad \tilde{C}r(h)\|u-E_hu\|_{a,\Omega}. \tag{3.4}$$

Note that (3.2) implies

$$u-E_hu=w^h-R_hw^h+u-w^h.$$

Hence we obtain (3.3) from (3.4). This completes the proof. □

3.1 A posteriori error estimators

Following the element residual $\tilde{\mathcal{R}}_T(u_{i,h})$ and the jump residual $\tilde{J}_E(u_{i,h})$ for (2.4), we now define an element residual $\mathcal{R}_T(E_hu)$ and a jump residual $J_E(E_hu)$ for (2.10) as follows:

$$\begin{aligned}\mathcal{R}_T(E_hu) &= \lambda^hE_hu+\nabla\cdot(A\nabla E_hu)-cE_hu \quad \text{in } T\in\mathcal{T}_h,\\ J_E(E_hu) &= -A\nabla(E_hu)^+\cdot\nu^+-A\nabla(E_hu)^-\cdot\nu^-\\ &= [[A\nabla E_hu]]_E\cdot\nu_E \qquad \text{on } E\in\mathcal{E}_h,\end{aligned}$$

where E , ν^+ and ν^- are defined as those of section 2.1.

For $T\in\mathcal{T}_h$, we define the local error indicator $\eta_h(E_hu,T)$ by

$$\eta_h^2(E_hu,T)=h_T^2\|\mathcal{R}_T(E_hu)\|_{0,T}^2+\sum_{E\in\mathcal{E}_h,E\subset\partial T}\|J_E(E_hu)\|_{0,E}^2$$

and the oscillation $osc_h(E_hu,T)$ by

$$osc_h^2(E_hu,T)=h_T^2\|\mathcal{R}_T(E_hu)-\overline{\mathcal{R}_T(E_hu)}\|_{0,T}^2+\sum_{E\in\mathcal{E}_h,E\subset\partial T}h_E\|J_E(E_hu)-\overline{J_E(E_hu)}\|_{0,E}^2.$$

We define the error estimator $\eta_h(E_hu,\Omega)$ and the oscillation $osc_h(E_hu,\Omega)$ by

$$\eta_h^2(E_hu,\Omega)=\sum_{T\in\mathcal{T}_h,T\subset\Omega}\eta_h^2(E_hu,T)\quad\text{and}\quad osc_h^2(E_hu,\Omega)=\sum_{T\in\mathcal{T}_h,T\subset\Omega}osc_h^2(E_hu,T).$$

For any $U_h=(u_{h,1},\cdots,u_{h,q})\in V_h^q$, we let

$$\eta_h^2(U_h,\Omega)=\sum_{l=1}^{q}\eta_h^2(u_{h,l},\Omega)\quad\text{and}\quad osc_h^2(U_h,\Omega)=\sum_{l=1}^{q}osc_h^2(u_{h,l},\Omega)$$

and for any $E_hU=(E_hu_1,\cdots,E_hu_q)\in V_h^q$, we set

$$\eta_h^2(E_hU,\Omega)=\sum_{l=1}^{q}\eta_h^2(E_hu_l,\Omega)\quad\text{and}\quad osc_h^2(E_hU,\Omega)=\sum_{l=1}^{q}osc_h^2(E_hu_l,\Omega).$$

Given $h_0 \in (0,1)$, define

$$\tilde{r}(h_0) = \sup_{h \in (0,h_0)} r(h).$$

Theorem 3.2. *There exist constants C_1, C_2 and C_3, which only depend on the shape regularity constant γ^*, coercivity constant c_a and continuity constant C_a of the bilinear form, such that*

$$\|u - E_h u\|_{a,\Omega} \le C_1 \eta_h(E_h u, \Omega) \tag{3.5}$$

and

$$C_2^2 \eta_h^2(E_h u, \Omega) - C_3^2 osc_h^2(E_h u, \Omega) \le \|u - E_h u\|_{a,\Omega}^2 \tag{3.6}$$

provided $h_0 \ll 1$. Consequently,

$$|\lambda - \lambda^h| \lesssim \eta_h^2(E_h u, \Omega)$$

and

$$\eta_h^2(E_h u, \Omega) - osc_h^2(E_h u, \Omega) \lesssim |\lambda - \lambda^h|.$$

Proof. Recall that $Lw^h = \sum_{l=1}^{q} \alpha_{h,l}(u)\lambda_{h,l} K u_{h,l}$. We obtain from (2.6) and (2.7) that

$$\|w^h - R_h w^h\|_{a,\Omega} \le \tilde{C}_1 \eta_h(E_h u, \Omega) \tag{3.7}$$

and

$$\tilde{C}_2^2 \eta_h^2(E_h u, \Omega) - \tilde{C}_3^2 osc_h^2(E_h u, \Omega) \le \|w^h - R_h w^h\|_{a,\Omega}^2. \tag{3.8}$$

Combining (3.2), (3.3), (3.7) with (3.8), we complete the proof. In particular, we may choose the constants C_1, C_2 and C_3 satisfying

$$C_1 = \tilde{C}_1(1 + \tilde{C}\tilde{r}(h_0)), \quad C_2 = \tilde{C}_2(1 - \tilde{C}\tilde{r}(h_0)), \quad C_3 = \tilde{C}_3(1 - \tilde{C}\tilde{r}(h_0)). \tag{3.9}$$

$\square$

3.2 Adaptive algorithm

Recall that the adaptive procedure consists of loops of the form

$$\boxed{\textbf{Solve} \rightarrow \textbf{Estimate} \rightarrow \textbf{Mark} \rightarrow \textbf{Refine}}$$

We assume that the solutions of the finite dimensional problems can be solved to any accuracy efficiently.[1] The a posteriori error estimators are an essential part of the **Estimate** step. In the following discussion, we use $\eta_h(U_h, \Omega)$ defined above as the a posteriori error estimator.

---

[1] In fact, we have ignored two important practical issues: the inexact solution of the resulting algebraic system and the numerical integration. We remark the discussion about the inexact solution in Section 7.

Algorithm C

Choose a parameter $0 < \theta < 1$.

1. Pick an initial mesh $\mathcal{T}_{h_0}$ with mesh size h_0.
2. Solve the system on $\mathcal{T}_{h_0}$ for the discrete solutions $(\lambda_{h_0,l}, u_{h_0,l})(l = 1, \cdots, q)$.
3. Let $k = 0$.
4. Compute the local indicators $\eta_{h_k}(u_{h_{k,l}}, T)(l = 1, \cdots, q)$.
5. Construct $\mathcal{M}_{h_k} \subset \mathcal{T}_{h_k}$ by **Marking Strategy** E and parameter θ.
6. Refine $\mathcal{T}_{h_k}$ to get a new conforming mesh $\mathcal{T}_{h_{k+1}}$ by Procedure **REFINE**.
7. Solve the system on $\mathcal{T}_{h_{k+1}}$ for the discrete solution $(\lambda_{h_{k+1},l}, u_{h_{k+1},l})(l = 1, \cdots, q)$.
8. Let $k = k + 1$ and go to Step 4.

Our **Algorithm** C is nothing but **Algorithm** C_0 when **Marking Strategy** E_0 is replaced by the following **Marking Strategy** E.

Marking Strategy E

Given a parameter $0 < \theta < 1$.

1. Construct a minimal subset $\mathcal{M}_{h_k}$ of $\mathcal{T}_{h_k}$ by selecting some elements in $\mathcal{T}_{h_k}$ such that
$$\sum_{T \in \mathcal{M}_{h_k}} \eta^2_{h_k}(U_{h_k}, T) \geq \theta \eta^2_{h_k}(U_{h_k}, \Omega).$$
2. Mark all the elements in $\mathcal{M}_{h_k}$.

We shall now present the following property of eigenspace approximation that will play a crucial role in our analysis.

Lemma 3.1. *Given constant* $\theta \in (0, 1)$. *If*

$$\sum_{T \in \mathcal{M}_H} \eta^2_H(U_H, T) \geq \theta \eta^2_H(U_H, \Omega), \tag{3.10}$$

then for any orthonormal basis $\{u_l\}_{l=1}^q$ *of* $M(\lambda)$*, there holds*

$$\sum_{T\in\mathcal{M}_H}\eta_H^2(E_HU,T)\geq\theta'\eta_H^2(E_HU,\Omega),\tag{3.11}$$

where $E_HU=(E_Hu_1,\cdots,E_Hu_q)$ *and* $\theta'=\frac{\theta}{4q^2}\in(0,1)$.

Proof. We denote E_Hu_l as $v_{H,l}$. Since $M_H(\lambda)=\text{span}\{u_{H,1},\cdots,u_{H,q}\}$, we have that there exists q constants $\beta_{H,j}^l(j=1,\cdots,q)$ satisfying $\sum_{j=1}^q\left(\beta_{H,j}^l\right)^2=1$ such that

$$v_{H,l}=\sum_{j=1}^q\beta_{H,j}^lu_{H,j},\quad l=1,\cdots,q.$$

We may analyze as follows

$$\begin{aligned}\eta_H^2(E_HU,\Omega) &= \sum_{l=1}^q\eta_H^2(\sum_{j=1}^q\beta_{H,j}^lu_{H,j},\Omega)\\ &\leq 2\sum_{l=1}^q\sum_{j=1}^q\left(\beta_{H,j}^l\right)^2\eta_H^2(u_{H,j},\Omega)\\ &= 2\sum_{j=1}^q\sum_{l=1}^q\left(\beta_{H,j}^l\right)^2\eta_H^2(u_{H,j},\Omega),\end{aligned}$$

which together with the fact that $|\beta_{H,j}^l|\leq1$ yields

$$\eta_H^2(E_HU,\Omega)\leq 2q\eta_H^2(U_H,\Omega).\tag{3.12}$$

We see from the definition of E_H that $\{v_{H,l}\}_{l=1}^q$ is an orthonormal basis of $M_H(\lambda)$, namely, $M_H(\lambda)=\text{span}\{u_{H,1},\cdots,u_{H,q}\}=\text{span}\{v_{H,1},\cdots,v_{H,q}\}$. So there exist q constants $\hat\beta_{H,j}^l(j=1,\cdots,q)$ satisfying $\sum_{j=1}^q\left(\hat\beta_{H,j}^l\right)^2=1$ such that

$$u_{H,l}=\sum_{j=1}^q\hat\beta_{H,j}^lv_{H,j},\quad l=1,\cdots,q.$$

Similarly, from the following estimation

$$\begin{aligned}\sum_{T\in\mathcal{M}_H}\eta_H^2(U_H,T) &= \sum_{l=1}^q\sum_{T\in\mathcal{M}_H}\eta_H^2(\sum_{j=1}^q\hat\beta_{H,j}^lv_{H,j},T)\\ &\leq 2\sum_{l=1}^q\sum_{j=1}^q\left(\hat\beta_{H,j}^l\right)^2\sum_{T\in\mathcal{M}_H}\eta_H^2(v_{H,j},T)\\ &= 2\sum_{j=1}^q\sum_{l=1}^q\left(\hat\beta_{H,j}^l\right)^2\sum_{T\in\mathcal{M}_H}\eta_H^2(v_{H,j},T),\end{aligned}$$

we obtain

$$\sum_{T\in\mathcal{M}_H}\eta_H^2(U_H,T)\leq 2q\sum_{T\in\mathcal{M}_H}\eta_H^2(E_HU,T).\tag{3.13}$$

Combing (3.10), (3.12) and (3.13), we arrive at

$$\begin{aligned}\eta_H^2(E_HU,\Omega) &\le 2q\eta_H^2(U_H,\Omega)\\ &\le 2q\frac{1}{\theta}\sum_{T\in\mathcal{M}_H}\eta_H^2(U_H,T)\\ &\le 4q^2\frac{1}{\theta}\sum_{T\in\mathcal{M}_H}\eta_H^2(E_HU,T),\end{aligned}$$

which is nothing but (3.11) with $\theta' = \frac{\theta}{4q^2}$. $\square$

4 Convergence

Following Theorem 3.1, we can establish some relationship between the two level approximations, which will be used in our analysis for both convergence and quasi-optimal complexity.

Lemma 4.1. *Let $h, H \in (0, h_0)$, $\{u_l\}_{l=1}^q$ be any orthonormal basis of $M(\lambda)$, $\lambda^{H,l} = a(E_Hu_l, E_Hu_l)$, $w^{H,l} = \sum_{i=1}^q \alpha_{H,i}(u_l)\lambda_{H,i}Ku_{H,i}$ and $W^H \equiv (w^{H,1}\cdots, w^{H,q})$. Then*

$$\|U - E_hU\|_{a,\Omega} = \|W^H - R_hW^H\|_{a,\Omega} + \mathcal{O}(\tilde r(h_0))\left(\|U - E_hU\|_{a,\Omega} + \|U - E_HU\|_{a,\Omega}\right), \tag{4.1}$$

$$osc_h(E_hU,\Omega) = \widetilde{osc}_h(R_hW^H,\Omega) + \mathcal{O}(\tilde r(h_0))\left(\|U - E_hU\|_{a,\Omega} + \|U - E_HU\|_{a,\Omega}\right), \tag{4.2}$$

and

$$\eta_h(E_hU,\Omega) = \tilde\eta_h(R_hW^H,\Omega) + \mathcal{O}(\tilde r(h_0))\left(\|U - E_hU\|_{a,\Omega} + \|U - E_HU\|_{a,\Omega}\right). \tag{4.3}$$

Proof. It is sufficient to prove that for any $u \in M(\lambda)$, $w^h = \sum_{l=1}^q \alpha_{h,l}(u)\lambda_{h,l}Ku_{h,l}$, and $w^H = \sum_{l=1}^q \alpha_{H,l}(u)\lambda_{H,l}Ku_{H,l}$, the following equalities hold,

$$\|u - E_hu\|_{a,\Omega} = \|w^H - R_hw^H\|_{a,\Omega} + \mathcal{O}(\tilde r(h_0))\left(\|u - E_hu\|_{a,\Omega} + \|u - E_Hu\|_{a,\Omega}\right), \tag{4.4}$$

$$osc_h(E_hu,\Omega) = \widetilde{osc}_h(R_hw^H,\Omega) + \mathcal{O}(\tilde r(h_0))\left(\|u - E_hu\|_{a,\Omega} + \|u - E_Hu\|_{a,\Omega}\right) \tag{4.5}$$

and

$$\eta_h(E_hu,\Omega) = \tilde\eta_h(R_hw^H,\Omega) + \mathcal{O}(\tilde r(h_0))\left(\|u - E_hu\|_{a,\Omega} + \|u - E_Hu\|_{a,\Omega}\right). \tag{4.6}$$

First, we prove (4.4). We see that

$$\begin{aligned}\|R_h(w^h - w^H) + w^H - u\|_{a,\Omega} &\lesssim \|w^h - w^H\|_{a,\Omega} + \|u - w^H\|_{a,\Omega}\\ &\lesssim \|u - w^H\|_{a,\Omega} + \|u - w^h\|_{a,\Omega},\end{aligned}$$

which together with (3.4) leads to

$$\|R_h(w^h - w^H) + w^H - u\|_{a,\Omega} \lesssim \tilde r(h_0)(\|u - E_Hu\|_{a,\Omega} + \|u - E_hu\|_{a,\Omega}). \tag{4.7}$$

Note that (3.2) implies

$$u - E_hu = w^H - R_hw^H + R_h(w^H - w^h) + u - w^H, \tag{4.8}$$

we get (4.4) from (4.7).

Next, we prove (4.5). We obtain from Lemma 2.1 that

$$\widetilde{osc}_h(R_h(w^h - w^H), \Omega) \lesssim \|R_h(w^H - w^h)\|_{a,\Omega}, \tag{4.9}$$

which together with (3.4) and (4.7) yields

$$\widetilde{osc}_h(R_h(w^H - w^h), \Omega) \lesssim \tilde{r}(h_0)(\|u - E_H u\|_{a,\Omega} + \|u - E_h u\|_{a,\Omega}). \tag{4.10}$$

Due to $E_h u = R_h w^H + R_h(w^h - w^H)$, we conclude from the definition of oscillation that

$$\widetilde{osc}_h(R_h w^h, \Omega) = \widetilde{osc}_h(R_h w^H + R_h(w^h - w^H), \Omega),$$

then we arrive at (4.5), together with (4.10).

We now prove (4.6). By (2.7), we have

$$\begin{aligned} \tilde{\eta}_h(R_h(w^h - w^H), \Omega) &\lesssim \|(w^h - w^H) - R_h(w^h - w^H)\|_{a,\Omega} + \widetilde{osc}_h(R_h(w^h - w^H), \Omega) \\ &\lesssim \|u - w^h\|_{a,\Omega} + \|u - w^H\|_{a,\Omega} + \|R_h(w^h - w^H)\|_{a,\Omega}, \end{aligned}$$

where (4.9) is used in the last inequality. Using (3.4) and (4.7), we obtain

$$\tilde{\eta}_h(R_h(w^h - w^H), \Omega) \quad \lesssim \quad \tilde{r}(h_0)(\|u - E_H u\|_{a,\Omega} + \|u - E_h u\|_{a,\Omega}). \tag{4.11}$$

From (4.11) and the fact that

$$\tilde{\eta}_h(R_h w^h, \Omega) = \tilde{\eta}_h(R_h w^H + R_h(w^h - w^H), \Omega),$$

we get

$$\tilde{\eta}_h(R_h w^h, \Omega) = \tilde{\eta}_h(R_h w^H, \Omega) + \mathcal{O}(\tilde{r}(h_0)) \left(\|u - E_h u\|_{a,\Omega} + \|u - E_H u\|_{a,\Omega}\right),$$

which is nothing but (4.6) since $\tilde{\eta}_h(R_h w^h, \Omega) = \eta_h(E_h u, \Omega)$. □

We are now in the position to present and analyze the error reduction result.

Theorem 4.1. *Let $\lambda \in \mathbb{R}$ be some eigenpair of (2.9) with multiplicity q, $\{u_l\}_{l=1}^q$ be an orthonormal basis of $M(\lambda)$, and $\{(\lambda_{h_k,l}, u_{h_k,l}), l = 1, \cdots, q\}_{k \in \mathbb{N}_0}$ be a sequence of finite element solutions produced by* **Algorithm** *C of Section 3. Then there exist constants $\gamma > 0$ and $\alpha \in (0, 1)$, depending only on the shape regularity of meshes, C_a and c_a, the parameter used by* **Algorithm** *C, such that for any two consecutive iterates k and $k + 1$, we have*

$$\|U - E_{h_{k+1}} U\|_{a,\Omega}^2 + \gamma \eta_{h_{k+1}}^2 (E_{h_{k+1}} U, \Omega) \le \alpha^2 \left(\|U - E_{h_k} U\|_{a,\Omega}^2 + \gamma \eta_{h_k}^2 (E_{h_k} U, \Omega)\right) \tag{4.12}$$

provided $h_0 \ll 1$. Therefore, **Algorithm C** *converges with a linear rate α, namely, the n-th iterate solution $(\lambda^{h_n,l}, E_{h_n} u_l) (l = 1, \cdots, q)$ of* **Algorithm** *C satisfies*

$$\|U - E_{h_n} U\|_{a,\Omega}^2 + \gamma \eta_{h_n}^2 (E_{h_n} U, \Omega) \quad \le C_0 \alpha^{2n} \tag{4.13}$$

and

$$\lambda^{h_n,l} - \lambda \lesssim \alpha^{2n}, \tag{4.14}$$

where $C_0 = \|U - E_{h_0} U\|_{a,\Omega}^2 + \gamma \eta_{h_0}^2 (E_{h_0} U, \Omega)$.

Proof. For convenience, we use $(\lambda_{h,l}, u_{h,l})$, $(\lambda_{H,l}, u_{H,l})$ to denote $(\lambda_{h_{k+1},l}, u_{h_{k+1},l})$ and $(\lambda_{h_k,l}, u_{h_k,l})$, respectively. We see that it is sufficient to prove

$$\|U - E_hU\|_{a,\Omega}^2 + \gamma\eta_h^2(E_hU,\Omega) \le \alpha^2\big(\|U - E_HU\|_{a,\Omega}^2 + \gamma\eta_H^2(E_HU,\Omega)\big).$$

We derive from Lemma 3.1 that **Marking Strategy E** implies that there exists a constant $\theta' = \frac{\theta}{4q^2} \in (0,1)$ such that

$$\sum_{T\in\mathcal{M}_H} \eta_H^2(E_HU,T) \ge \theta'\eta_H^2(E_HU,\Omega). \tag{4.15}$$

Recall that $w^{h,l} = \sum_{i=1}^q \alpha_{h,i}(u_l)\lambda_{h,i}Ku_{h,i}$, $w^{H,l} = \sum_{i=1}^q \alpha_{H,i}(u_l)\lambda_{H,i}Ku_{H,i}$, we get from (4.15) that for $W^H \equiv (w^{H,1},\cdots,w^{H,q})$, **Marking Strategy** E_0 is satisfied with $\theta = \theta'$. So, we conclude from Theorem 2.1 that there exist constants $\tilde{\gamma} > 0$ and $\xi \in (0,1)$ satisfying

$$\|W^H - R_hW^H\|_{a,\Omega}^2 + \tilde{\gamma}\tilde{\eta}_h^2(R_hW^H,\Omega) \le \xi^2\|W^H - E_HU\|_{a,\Omega}^2 + \tilde{\gamma}\eta_H^2(E_HU,\Omega), \tag{4.16}$$

where (3.2) is used when h is replaced by H.

By Lemma 4.1 and (3.4), we have that there exist constants $\hat{C}_1 > 0, \hat{C}_2 > 0$ such that

$$\begin{aligned}\|U - E_HU\|_{a,\Omega}^2 + \tilde{\gamma}\eta_H^2(E_HU,\Omega) &= \|W^H - E_HU\|_{a,\Omega}^2 + \tilde{\gamma}\eta_H^2(E_HU,\Omega) \\ &\quad + \hat{C}_1\tilde{r}^2(h_0)\|U - E_HU\|_{a,\Omega}^2 \end{aligned}\tag{4.17}$$

and

$$\begin{aligned}\|U - E_hU\|_{a,\Omega}^2 + \tilde{\gamma}\eta_h^2(E_hU,\Omega) &= \|W^H - R_hW^H\|_{a,\Omega}^2 + \tilde{\gamma}\tilde{\eta}_h^2(R_hW^H,\Omega) \\ &\quad + \hat{C}_2\tilde{r}^2(h_0)\big(\|U - E_hU\|_{a,\Omega}^2 + \|U - E_HU\|_{a,\Omega}^2\big). \end{aligned}\tag{4.18}$$

Combing (4.16), (4.17) with (4.18), we get when $h_0 \ll 1$ that

$$\begin{aligned} &\|U - E_hU\|_{a,\Omega}^2 + \frac{\tilde{\gamma}}{1 - \hat{C}_2\tilde{\gamma}^2(h_0)}\eta_h^2(E_hU,\Omega) \\ \le\ & \frac{\xi^2(1 - \hat{C}_1\tilde{\gamma}^2(h_0)) + \hat{C}_2\tilde{\gamma}^2(h_0)}{1 - \hat{C}_2\tilde{\gamma}^2(h_0)}\|U - E_HU\|_{a,\Omega}^2 + \frac{\xi^2}{1 - \hat{C}_2\tilde{\gamma}^2(h_0)}\tilde{\gamma}\eta_H^2(E_HU,\Omega). \end{aligned}$$

Since $h_0 \ll 1$ implies $\tilde{r}(h_0) \ll 1$, there exists $\delta_1 \in (0,1)$ such that $|\hat{C}_1\tilde{\gamma}^2(h_0)| \le \delta_1$ if $h_0 \ll 1$. Hence,

$$\begin{aligned} &\|U - E_hU\|_{a,\Omega}^2 + \frac{\tilde{\gamma}}{1 - \hat{C}_2\tilde{\gamma}^2(h_0)}\eta_h^2(E_hU,\Omega) \\ \le\ & \frac{\xi^2(1+\delta_1) + \hat{C}_2\tilde{\gamma}^2(h_0)}{1 - \hat{C}_2\tilde{\gamma}^2(h_0)}\left(\|U - E_HU\|_{a,\Omega}^2 + \frac{\xi^2\tilde{\gamma}}{\xi^2(1+\delta_1) + \hat{C}_2\tilde{\gamma}^2(h_0)}\eta_H^2(E_HU,\Omega)\right) \end{aligned}$$

when $h_0 \ll 1$. Note that the constant α defined by

$$\alpha = \left(\frac{\xi^2(1+\delta_1) + \hat{C}_2\tilde{\gamma}^2(h_0)}{1 - \hat{C}_2\tilde{\gamma}^2(h_0)}\right)^{1/2}$$

satisfies $\alpha \in (0,1)$ if $h_0 \ll 1$. We finally arrive at (4.12) by using the fact that

$$\frac{\xi^2\tilde{\gamma}}{\xi^2(1+\delta_1) + \hat{C}_2\tilde{\gamma}^2(h_0)} < \gamma,$$

where

$$\gamma = \frac{\tilde{\gamma}}{1 - \hat{C}_2\tilde{\gamma}^2(h_0)}. \tag{4.19}$$

This completes the proof. □

5 Complexity

We introduce a function approximation class as follows

$$\mathcal{A}_\gamma^s = \{v \in H : |v|_{s,\gamma} < \infty\},$$

where $\gamma > 0$ is some constant,

$$|v|_{s,\gamma} = \sup_{\varepsilon>0} \varepsilon \inf_{\{\mathcal{T}\subset\mathcal{T}_{h_0}:\inf(\|v-v_{\mathcal{T}}\|_{1,\Omega}^2+(\gamma+1)osc_{\mathcal{T}}^2(v_{\mathcal{T}},\mathcal{T}))^{1/2}\le\varepsilon\}} \big(\#\mathcal{T} - \#\mathcal{T}_{h_0}\big)^s$$

and $\mathcal{T} \subset \mathcal{T}_{h_0}$ means $\mathcal{T}$ is a refinement of $\mathcal{T}_{h_0}$. It is seen from the definition that, for all $\gamma > 0$, $\mathcal{A}_\gamma^s = \mathcal{A}_1^s$. For simplicity, here and hereafter, we use $\mathcal{A}^s$ to stand for $\mathcal{A}_1^s$, and use $|v|_s$ to denote $|v|_{s,\gamma}$. So $\mathcal{A}^s$ is the class of functions that can be approximated within a given tolerance ε by continuous piecewise polynomial functions over a partition $\mathcal{T}$ with number of degrees of freedom $\#\mathcal{T} - \#\mathcal{T}_{h_0} \le \varepsilon^{-1/s}|v|_s^{1/s}$.

To give the proof of optimal complexity of **Algorithm** C for the eigenvalue problem (2.9), we need some preparations. We know that in each mesh $\mathcal{T}_{h_k}$, $w^{h_k,l} = \sum_{i=1}^q \alpha_{h_k,i}(u_l)\lambda_{h_k,i}Ku_{h_k,i}$ satisfies

$$a(w^{h_k,l}, v) = (\sum_{i=1}^q \alpha_{h_k,i}(u_l)\lambda_{h_k,i}u_{h_k,i}, v) \quad \forall v \in V_{h_k}, \quad l = 1, \cdots, q. \tag{5.1}$$

Thanks to Theorem 3.1 and Lemma 4.1 and their proofs, we are able to analyze the complexity of adaptive finite element method for multiple eigenvalue problems by using the complexity result for boundary value problems, which is similar to what was demonstrated in the convergence analysis.

Using the similar procedure as in the proof of Theorem 4.1, we have

Lemma 5.1. *Let $\lambda \in \mathbb{R}$ be some eigenvalue of (2.9) with multiplicity q, and $\{u_l\}_{l=1}^q$ be an orthonormal basis of $M(\lambda)$. Let $(\lambda_{h_k,l}, u_{h_k,l}) \in \mathbb{R} \times V_{h_k}$ and $(\lambda_{h_{k+1},l}, u_{h_{k+1},l}) \in \mathbb{R} \times V_{h_{k+1}}$ $(l = 1, \cdots, q)$ be discrete solutions of (2.10) over a conforming mesh $\mathcal{T}_{h_k}$ and its refinement $\mathcal{T}_{h_{k+1}}$ with marked element $\mathcal{M}_{h_k}$. Suppose they satisfy the following property*

$$\begin{aligned} & \|U - E_{h_{k+1}}U\|_{a,\Omega}^2 + \gamma_* osc_{h_{k+1}}^2(E_{h_{k+1}}U, \Omega) \\ \le\ & \beta_*^2\Big(\|U - E_{h_k}U\|_{a,\Omega}^2 + \gamma_* osc_{h_k}^2(E_{h_k}U, \Omega)\Big), \end{aligned}$$

where $\gamma_* > 0, \beta_* > 0$ are some constants. Then for the associated boundary value problem (5.1), we have

$$\begin{aligned} & \|W^{h_k} - R_{h_{k+1}}W^{h_k}\|_{a,\Omega}^2 + \tilde{\gamma}_* \widetilde{osc}_{h_{k+1}}^2(R_{h_{k+1}}W^{h_k}, \Omega) \\ \le\ & \tilde{\beta}_*^2\Big(\|W^{h_k} - R_{h_k}W^{h_k}\|_{a,\Omega}^2 + \tilde{\gamma}_* \widetilde{osc}_{h_k}^2(E_{h_k}U, \Omega)\Big) \end{aligned}$$

with

$$\tilde{\beta}_* = \left(\frac{\beta_*^2(1+\delta_1) + \hat{C}_3\tilde{\gamma}^2(h_0)}{1 - \hat{C}_3\tilde{\gamma}^2(h_0)}\right)^{1/2}, \quad \tilde{\gamma}_* = \frac{\gamma_*}{1 - \hat{C}_3\tilde{\gamma}^2(h_0)}, \tag{5.2}$$

where C_5 is some constant depending on A , $\tilde{C}$ and C_*, $\delta_1 \in (0,1)$ is some constant as shown in the proof Theorem 4.1.

Proof. By Lemma 4.1 and (3.4), we have that there exists constant $\hat{C}_3 > 0$ such that

$$\begin{aligned}\|W^H - R_H W^H\|_{a,\Omega}^2 + \gamma_* \widetilde{osc}_H^2(R_H W^H, \Omega) &= \|U - E_H U\|_{a,\Omega}^2 + \gamma_* \widetilde{osc}_H^2(E_H U, \Omega) \\ &\quad + \hat{C}_1 \tilde{r}^2(h_0) \|W^H - R_H W^H\|_{a,\Omega}^2,\end{aligned}$$

and

$$\begin{aligned}\|W^H - R_h W^H\|_{a,\Omega}^2 + \gamma_* \widetilde{osc}_h^2(R_h W^H, \Omega) &= \|U - E_h U\|_{a,\Omega}^2 + \gamma_* osc_h^2(E_h U, \Omega) \\ &\quad + \hat{C}_3 \tilde{r}^2(h_0) \big(\|W^H - R_h W^H\|_{a,\Omega}^2 + \|W^H - R_H W^H\|_{a,\Omega}^2 \big).\end{aligned}$$

Proceed the same procedure as in the proof of Theorem 4.1, we get

$$\|W^H - R_h W^H\|_{a,\Omega}^2 + \tilde{\gamma}_* \widetilde{osc}_h^2(R_h W^H, \Omega) \le \tilde{\beta}_*^2 \big(\|W^H - R_H W^H\|_{a,\Omega}^2 + \tilde{\gamma}_* \widetilde{osc}_H^2(R_H W^H, \Omega) \big)$$

with

$$\tilde{\beta}_* = \left(\frac{\beta_*^2 (1 + \delta_1) + \hat{C}_3 \tilde{\gamma}^2(h_0)}{1 - \hat{C}_3 \tilde{\gamma}^2(h_0)} \right)^{1/2}, \quad \tilde{\gamma}_* = \frac{\gamma_*}{1 - \hat{C}_3 \tilde{\gamma}^2(h_0)},$$

where C_3 is some positive constant and $\delta_1 \in (0, 1)$ is some constant as shown in the proof Theorem 4.1. This completes the proof. $\square$

The following statement is a direct consequence of $E_{h_k} u_l = R_{h_k} w^{h_k, l}$, Proposition 2.2 and Lemma 5.1.

Corollary 5.1. *Let $\lambda \in \mathbb{R}$ be some eigenpair of (2.9) with multiplicity q, and $\{u_l\}_{l=1}^q$ be an orthonormal basis of $M(\lambda)$. Suppose that they satisfy the decrease property*

$$\begin{aligned}&\|U - E_{h_{k+1}} U\|_{a,\Omega}^2 + \gamma_* osc_{h_{k+1}}^2(E_{h_{k+1}} U, \Omega) \\ &\le \beta_*^2 \Big(\|U - E_{h_k} U\|_{a,\Omega}^2 + \gamma_* osc_{h_k}^2(E_{h_k} U, \Omega) \Big)\end{aligned}$$

with constants $\gamma_* > 0$ and $\beta_* \in (0, \sqrt{\frac{1}{2}})$. Then the set $\mathcal{R} = R_{\mathcal{T}_{h_k} \to \mathcal{T}_{h_{k+1}}}$ satisfies the following inequality

$$\sum_{T \in \mathcal{R}} \eta_{h_k}^2(E_{h_k} U, T) \ge \hat{\theta} \sum_{T \in \mathcal{T}_{h_k}} \eta_{h_k}^2(E_{h_k} U, T)$$

with $\hat{\theta} = \frac{\tilde{C}_2^2 (1 - 2\tilde{\beta}_*^2)}{\tilde{C}_0 (\tilde{C}_1^2 + (1 + 2C_*^2 \tilde{C}_1^2) \tilde{\gamma}_*)}$ and $\tilde{C}_0 = \max(1, \frac{\tilde{C}_3^2}{\tilde{\gamma}_*})$, where $\tilde{\beta}_*$ and $\tilde{\gamma}_*$ are defined in (5.2) with δ_2 being chosen such that ${\tilde{\beta}_*}^2 \in (0, \frac{1}{2})$.

Lemma 5.2. *Let $\lambda \in \mathbb{R}$ be some eigenvalue of (2.9) with multiplicity q, $\{u_l\}_{l=1}^q$ be an orthonormal basis of $M(\lambda)$, $u_l \in \mathcal{A}^s (l = 1, \cdots, q)$ and $\mathcal{T}_{h_k}$ be a conforming partition obtained from $\mathcal{T}_{h_0}$. Let $\mathcal{T}_{h_{k+1}}$ be a mesh created from $\mathcal{T}_{h_k}$ upon making the set $\mathcal{M}_{h_k}$ according with* **Marking Strategy** *E with $\theta \in (0, \frac{C_2^2 \gamma}{C_3^2 (C_1^2 + (1 + 2C_*^2 C_1^2) \gamma)})$. Let $\{(\lambda_{h_k, l}, u_{h_k, l}), l = 1, \cdots, q\}_{k \in \mathbb{N}_0}$ be discrete solutions of (2.10) over a conforming mesh $\mathcal{T}_{h_k}$ and $M_{h_k}(\lambda) = span\{u_{h_k, 1}, \cdots, u_{h_k, q}\}$. Then*

$$\#\mathcal{M}_{h_k} \le C \Big(\|U - E_{h_k} U\|_{a,\Omega}^2 + \gamma osc_{h_k}^2(E_{h_k} U, \Omega) \Big)^{-1/2s} \Big(q^{\frac{1}{2s} - 1} \sum_{l=1}^q |u_l|_s^{1/s} \Big), \tag{5.3}$$

where the constant C depends on the discrepancy between θ and $\frac{C_2^2 \gamma}{C_3^2 (C_1^2 + (1 + 2C_*^2 C_1^2) \gamma)}$.

Proof. Let $\beta, \beta_1 \in (0,1)$ satisfy $\beta_1 \in (0,\beta)$ and

$$\theta < \frac{C_2^2\gamma}{C_3^2(C_1^2 + (1+2C_*^2C_1^2)\gamma)}(1-\beta^2).$$

Choose

$$\varepsilon = \frac{1}{\sqrt{2}}\beta_1\Big(\|U - E_{h_k}U\|_a^2 + \gamma osc_{h_k}^2(E_{h_k}U,\Omega)\Big)^{1/2}$$

and let $\mathcal{T}_{h_\varepsilon}$ be a refinement of $\mathcal{T}_{h_0}$ with minimal degrees of freedom satisfying

$$\|u_l - E_{h_\varepsilon}u_l\|_{a,\Omega}^2 + (\gamma+1)osc_{h_\varepsilon}^2(E_{h_\varepsilon}u_l,\Omega) \le \frac{\varepsilon^2}{q}, \quad l = 1,\cdots,q,$$

which means

$$\|U - E_{h_\varepsilon}U\|_{a,\Omega}^2 + (\gamma+1)osc_{h_\varepsilon}^2(E_{h_\varepsilon}U,\Omega) \le \varepsilon^2. \tag{5.4}$$

We get from the definition of $\mathcal{A}^s$ that

$$\begin{aligned}&\#\mathcal{T}_{h_\varepsilon} - \#\mathcal{T}_{h_0}\\ \le\ & (\frac{1}{\sqrt{2}}\beta_1)^{-1/s}\Big(\|U - E_{h_k}U\|_{a,\Omega}^2 + \gamma osc_{h_k}^2(E_{h_k}U,\Omega)\Big)^{-1/2s} q^{1/2s}|u_l|_s^{1/s}, \quad l = 1,\cdots,q,\end{aligned}$$

which implies

$$\begin{aligned}&\#\mathcal{T}_{h_\varepsilon} - \#\mathcal{T}_{h_0}\\ \le\ & (\frac{1}{\sqrt{2}}\beta_1)^{-1/s}\Big(\|U - E_{h_k}U\|_{a,\Omega}^2 + \gamma osc_{h_k}^2(E_{h_k}U,\Omega)\Big)^{-1/2s}\Big(q^{\frac{1}{2s}-1}\sum_{l=1}^q |u_l|_s^{1/s}\Big).\end{aligned}$$

Let $\mathcal{T}_{h_{k,+}}$ be the smallest common refinement of $\mathcal{T}_{h_k}$ and $\mathcal{T}_{h_\varepsilon}$. Note that both $\mathcal{T}_{h_k}$ and $\mathcal{T}_{h_\varepsilon}$ are refinements of $\mathcal{T}_{h_0}$, we have that the number of elements in $\mathcal{T}_{h_{k,+}}$ that are not in $\mathcal{T}_{h_k}$ is less than the number of elements that must be added to go from $\mathcal{T}_{h_0}$ to $\mathcal{T}_{h_\varepsilon}$, namely,

$$\#\mathcal{T}_{h_{k,+}} - \#\mathcal{T}_{h_k} \le \#\mathcal{T}_{h_\varepsilon} - \#\mathcal{T}_{h_0}.$$

Let $w^{h_\varepsilon,l} = \sum_{i=1}^q \alpha_{h_\varepsilon,i}(u_l)\lambda_{h_\varepsilon,i}Ku_{h_\varepsilon,i} = K\big(\sum_{i=1}^q \alpha_{h_\varepsilon,i}(u_l)\lambda_{h_\varepsilon,i}u_{h_\varepsilon,i}\big)$ $(l = 1,\cdots q)$, namely

$$Lw^{h_\varepsilon,l} = \sum_{i=1}^q \alpha_{h_\varepsilon,i}(u_l)\lambda_{h_\varepsilon,i}u_{h_\varepsilon,i}.$$

We obtain from Lemma 2.1 and the Young inequality that

$$\widetilde{osc}_{h_{k,+}}^2(R_{h_{k,+}}W^{h_\varepsilon},\Omega) \ \le\ 2\widetilde{osc}_{h_\varepsilon}^2(R_{h_\varepsilon}W^{h_\varepsilon},\Omega) + 2C_*^2\|R_{h_{k,+}}W^{h_\varepsilon} - R_{h_\varepsilon}W^{h_\varepsilon}\|_{a,\Omega}^2.$$

Note that $\mathcal{T}_{h_{k,+}}$ is a refinement of $\mathcal{T}_{h_\varepsilon}$, L^2-projection error are monotone and the following orthogonality

$$\|W^{h_\varepsilon} - R_{h_{k,+}}W^{h_\varepsilon}\|_{a,\Omega}^2 = \|W^{h_\varepsilon} - R_{h_\varepsilon}W^{h_\varepsilon}\|_{a,\Omega}^2 - \|R_{h_{k,+}}W^{h_\varepsilon} - R_{h_\varepsilon}W^{h_\varepsilon}\|_{a,\Omega}^2$$

is valid, we arrive at

$$\begin{aligned}&\|W^{h_\varepsilon} - R_{h_{k,+}}W^{h_\varepsilon}\|_{a,\Omega}^2 + \frac{1}{2C_*^2}\widetilde{osc}_{h_{k,+}}^2(R_{h_{k,+}}W^{h_\varepsilon},\Omega)\\ \le\ & \|W^{h_\varepsilon} - R_{h_\varepsilon}W^{h_\varepsilon}\|_{a,\Omega}^2 + \frac{1}{C_*^2}osc_{h_\varepsilon}^2(R_{h_\varepsilon}W^{h_\varepsilon},\Omega).\end{aligned}$$

Since (2.8) implies $\tilde{\gamma} \leq \frac{1}{2C_*^2}$, we obtain that $\sigma \equiv \frac{1}{C_*^2} - \tilde{\gamma} \in (0,1)$ and

$$\begin{aligned} & \|W^{h_\varepsilon} - R_{h_{k,+}} W^{h_\varepsilon}\|_{a,\Omega}^2 + \tilde{\gamma}\widetilde{osc}_{h_{k,+}}^2(R_{h_{k,+}} W^{h_\varepsilon}, \Omega) \\ \leq \quad & \|W^{h_\varepsilon} - R_{h_\varepsilon} W^{h_\varepsilon}\|_{a,\Omega}^2 + \frac{1}{C_*^2} osc_{h_\varepsilon}^2(R_{h_\varepsilon} W^{h_\varepsilon}, \Omega) \\ \leq \quad & \|W^{h_\varepsilon} - R_{h_\varepsilon} W^{h_\varepsilon}\|_{a,\Omega}^2 + (\tilde{\gamma} + \sigma) osc_{h_\varepsilon}^2(R_{h_\varepsilon} W^{h_\varepsilon}, \Omega). \end{aligned}$$

Applying the similar argument in the proof of Theorem 4.1, we may conclude that

$$\begin{aligned} & \|U - E_{h_{k,+}} U\|_{a,\Omega}^2 + \gamma osc_{h_{k,+}}^2(E_{h_{k,+}} U, \Omega) \\ \leq \quad & \beta_0^2 \Big(\|U - E_{h_\varepsilon} U\|_{a,\Omega}^2 + (\gamma + \sigma) osc_{h_\varepsilon}^2(E_{h_\varepsilon} U, \Omega) \Big) \\ \leq \quad & \beta_0^2 \Big(\|U - E_{h_\varepsilon} U\|_{a,\Omega}^2 + (\gamma + 1) osc_{h_\varepsilon}^2(E_{h_\varepsilon} U, \Omega) \Big), \end{aligned} \tag{5.5}$$

where

$$\beta_0 = \left(\frac{1 + \delta_1 + \hat{C}_2 \tilde{\gamma}^2(h_0)}{1 - \hat{C}_2 \tilde{\gamma}^2(h_0)} \right)^{1/2},$$

and δ_1 is the constant appearing in the proof of Theorem 4.1. Combining (5.4) and (5.5), we then arrive at

$$\|U - E_{h_{k,+}} U\|_{a,\Omega}^2 + \gamma osc_{h_{k,+}}^2(E_{h_{k,+}} U, \Omega) \leq \breve{\beta}^2 \Big(\|U - E_{h_k} U\|_a^2 + \gamma osc_{h_k}^2(E_{h_k} U, \Omega) \Big)$$

with $\breve{\beta} = \frac{1}{\sqrt{2}} \beta_0 \beta_1$.

Let $\delta_1 \in (0,1)$ be some constant satisfying

$$(1 + \delta_1)^2 \beta_1^2 \leq \beta^2, \tag{5.6}$$

which implies

$$(1 + \delta_1) \beta_1^2 < 1. \tag{5.7}$$

We see from $h_0 \ll 1$ and (5.7) that $\breve{\beta}^2 \in (0, \frac{1}{2})$. Thus we get from Corollary 5.1 that $\mathcal{T}_{h_{k,+}}$ satisfies

$$\sum_{T \in \mathcal{M}_{h_{k,+}}} \eta_{h_k}^2(E_{h_k} U, T) \geq \breve{\theta} \sum_{T \in \mathcal{T}_{h_k}} \eta_{h_k}^2(E_{h_k} U, T),$$

where $\breve{\theta} = \frac{\tilde{C}_2^2(1 - 2\hat{\beta}^2)}{\tilde{C}_0(\tilde{C}_1^2 + (1 + 2C_*^2 \tilde{C}_1^2)\hat{\gamma})}$, $\quad \hat{\gamma} = \frac{\gamma}{1 - \hat{C}_3 \tilde{\gamma}^2(h_0)}$, $\tilde{C}_0 = max(1, \frac{\tilde{C}_3^2}{\hat{r}})$ and

$$\hat{\beta} = \left(\frac{\breve{\beta}^2(1 + \delta_1) + \hat{C}_3 \tilde{\gamma}^2(h_0)}{1 - \hat{C}_3 \tilde{\gamma}^2(h_0)} \right)^{1/2}.$$

From the definition of γ (see (4.19)) and $\tilde{\gamma}$ (see (2.8)), we obtain that $\hat{\gamma} < 1$. On the other hand, we have $\tilde{C}_3 > 1$ and hence $\tilde{C}_0 = \frac{\tilde{C}_3^2}{\hat{\gamma}}$. Consequently, we can write $\breve{\theta}$ as $\breve{\theta} = \frac{\tilde{C}_2^2(1 - 2\hat{\beta}^2)}{\tilde{C}_3^2(\frac{\tilde{C}_1^2}{\hat{\gamma}} + (1 + 2C_*^2 \tilde{C}_1^2))}$.

Since $h_0 \ll 1$, we obtain that $\hat{\gamma} > \gamma$ and $\hat{\beta} \in (0, \frac{1}{\sqrt{2}}\beta)$ from (5.6). Using (3.9), we get that

$$\begin{aligned} \breve{\theta} = \frac{\tilde{C}_2^2(1 - 2\hat{\beta}^2)}{\tilde{C}_3^2(\frac{\tilde{C}_1^2}{\hat{\gamma}} + (1 + 2C_*^2 \tilde{C}_1^2))} \geq \frac{\tilde{C}_2^2}{\tilde{C}_3^2(\frac{\tilde{C}_1^2}{\hat{\gamma}} + (1 + 2C_*^2 \tilde{C}_1^2))}(1 - \beta^2) \\ = \quad \frac{\frac{C_2^2}{(1 - \tilde{C}\tilde{r}(h_0))^2}}{\frac{C_3^2}{(1 - \tilde{C}\tilde{r}(h_0))^2}\big(\frac{C_1^2}{(1 + \tilde{C}\tilde{r}(h_0))^2 \hat{\gamma}} + (1 + 2C_*^2 \frac{C_1^2}{(1 + \tilde{C}\tilde{r}(h_0))^2})\big)}(1 - \beta^2), \end{aligned}$$

which together with the fact that $h_0 \ll 1$ and $\hat{\gamma} > \gamma$ yields

$$\check{\theta} \geq \frac{C_2^2}{C_3^2(\frac{C_1^2}{\gamma} + (1+2C_*^2C_1^2))}(1-\beta^2) = \frac{C_2^2\gamma}{C_3^2(C_1^2 + (1+2C_*^2C_1^2)\gamma)}(1-\beta^2) > \theta.$$

Therefore, we have

$$\begin{aligned}\#\mathcal{M}_{h_k} &\leq \#\mathcal{R} \leq \#\mathcal{T}_{h_{k,+}} - \#\mathcal{T}_{h_k} \leq \#\mathcal{T}_{h_\varepsilon} - \#\mathcal{T}_{h_0} \\ &\leq (\frac{1}{\sqrt{2}}\beta_1)^{-1/s}\Big(\|U - E_{h_k}U\|_{a,\Omega}^2 + \gamma osc_{h_k}^2(E_{h_k}U,\Omega)\Big)^{-1/2s}\Big(q^{\frac{1}{2s}-1}\sum_{l=1}^{q}|u_l|_s^{1/s}\Big).\end{aligned}$$

This is the desired estimate (5.3) with an explicit dependence on the discrepancy between θ and $\frac{C_2^2\gamma}{C_3^2(C_1^2+(1+2C_*^2C_1^2)\gamma)}$ via β_1. This completes the proof. $\square$

We are now ready to show that **Algorithm** C possesses quasi-optimal complexity.

Theorem 5.1. *Let $\lambda \in \mathbb{R}$ be some eigenvalue of (2.9) with multiplicity q, $\{u_l\}_{l=1}^q$ be an orthonormal basis of $M(\lambda)$, and $u_l \in \mathcal{A}^s (l = 1, \cdots, q)$. Let $\{(\lambda_{h_k,l}, u_{h_k,l}), l = 1, \cdots, q\}_{k\in\mathbb{N}_0}$ be a sequence of finite element solutions produced by* **Algorithm** *C of Section 3 and $M_{h_k}(\lambda) = span\{u_{h_k,1}, \cdots, u_{h_k,q}\}$. Then the n-th iterate solution space $M_{h_n}(\lambda)$ of* **Algorithm** *C satisfies the quasi-optimal bound*

$$\|U - E_{h_n}U\|_{a,\Omega}^2 + \gamma osc_{h_n}^2(E_{h_n}U,\Omega) \lesssim (\#\mathcal{T}_{h_n} - \#\mathcal{T}_{h_0})^{-2s},$$
$$\lambda^{h_n,l} - \lambda \lesssim (\#\mathcal{T}_{h_n} - \#\mathcal{T}_{h_0})^{-2s},$$

where the hidden constant depends on the exact solution $(\lambda, M(\lambda))$ and the discrepancy between θ and $\frac{C_2^2\gamma}{C_3^2(C_1^2+(1+2C_*^2C_1^2)\gamma)}$.

Proof. We see from (3.6) that

$$\|U - E_{h_k}U\|_{a,\Omega}^2 + \gamma\eta_{h_k}^2(E_{h_k}U,\Omega) \leq \check{C}\big(\|U - E_{h_k}U\|_{a,\Omega}^2 + \gamma osc_{h_k}^2(E_{h_k}U,\Omega)\big),$$

where $\check{C} = \max(1 + \frac{\gamma}{C_2^2}, \frac{C_3^2}{C_2^2})$. Hence, we get from (5.3) that

$$\#\mathcal{M}_{h_k} \leq (\frac{1}{\sqrt{2}}\beta_1)^{-1/s}\check{C}^{\frac{1}{2s}}\Big(\|U - E_{h_k}U\|_{a,\Omega}^2 + \gamma\eta_{h_k}^2(E_{h_k}U)\Big)^{-1/2s}\Big(q^{\frac{1}{2s}-1}\sum_{l=1}^{q}|u_l|_s^{1/s}\Big).$$

Note that Theorem 4.1 implies

$$\|U - E_{h_{k+1}}U\|_{a,\Omega}^2 + \gamma\eta_{h_{k+1}}^2(E_{h_{k+1}}U,\Omega) \leq \alpha^2\Big(\|U - E_{h_k}U\|_{a,\Omega}^2 + \gamma\eta_{h_k}^2(E_{h_k}U,\Omega)\Big).$$

Thus for $0 \leq k < n$, we arrive at

$$\begin{aligned}&\Big(\|U - E_{h_k}U\|_{a,\Omega}^2 + \gamma\eta_{h_k}^2(E_{h_k}U,\Omega)\Big)^{-1/2s} \\ \leq\ &\alpha^{(n-k)/s}\Big(\|U - E_{h_n}U\|_{a,\Omega}^2 + \gamma\eta_{h_n}^2(E_{h_n}u,\Omega)\Big)^{-1/2s}.\end{aligned}$$

We next employ Lemma 2.2 to deduce that

$$
\begin{aligned}
\#\mathcal{T}_{h_n} - \#\mathcal{T}_{h_0} &\lesssim \sum_{k=0}^{n-1} \#\mathcal{M}_{h_k} \\
\lesssim \quad & \Big(\sum_{l=1}^{q} |u_l|_s^{1/s}\Big) \sum_{k=0}^{n-1} \Big(\|U - E_{h_k}U\|_{a,\Omega}^2 + \gamma\eta_{h_k}^2(E_{h_k}U,\Omega)\Big)^{-1/2s} \\
\lesssim \quad & \Big(\|U - E_{h_n}U\|_{a,\Omega}^2 + \gamma\eta_{h_n}^2(E_{h_n}U,\Omega)\Big)^{-1/2s} \Big(\sum_{l=1}^{q} |u_l|_s^{1/s}\Big) \sum_{k=1}^{n} \alpha^{\frac{k}{s}},
\end{aligned}
$$

which together with the fact $\alpha < 1$ leads to

$$\#\mathcal{T}_{h_n} - \#\mathcal{T}_{h_0} \lesssim \Big(\|U - E_{h_n}U\|_{a,\Omega}^2 + \gamma\eta_{h_n}^2(E_{h_n}U,\Omega)\Big)^{-1/2s} \Big(\sum_{l=1}^{q} |u_l|_s^{1/s}\Big).$$

Since $osc(E_{h_n}U,\Omega) \le \eta_{h_n}(E_{h_n}U,\Omega)$, we conclude

$$\#\mathcal{T}_{h_n} - \#\mathcal{T}_{h_0} \lesssim \Big(\|U - E_{h_n}U\|_{a,\Omega}^2 + \gamma osc_{h_n}^2(E_{h_n}U,\Omega)\Big)^{-1/2s} \Big(\sum_{l=1}^{q} |u_l|_s^{1/s}\Big).$$

This completes the proof. □

6 Numerical examples

In this section, we show some numerical experiments for both linear finite elements and quadratic finite elements in three dimensions to illustrate the theoretical results obtained in this paper.

Our numerical experiments were carried out on LSSC-III in the State Key Laboratory of Scientific and Engineering Computing, Chinese Academy of Sciences, and our codes were based on the toolbox PHG of the State Key Laboratory of Scientific and Engineering Computing, Chinese Academy of Sciences.

For the convenience of analysis for our numerical results below, we define the following notations. Let $\lambda \in \mathbb{R}$ be some exact eigenvalue with multiplicity q, $M(\lambda)$ be the corresponding eigenfunction space as defined before, and $\{u_l\}_{l=1}^q$ be some orthonormal basis of $M(\lambda)$. Let $(\lambda_{h,l}, u_{h,l})(l = 1, \cdots, q)$ be the finite element approximation, $M_h(\lambda) = \text{span}\{u_{h,1}, \cdots, u_{h,q}\}$. For $M(\lambda)$ and $M_h(\lambda)$, we define the a posteriori error estimators $\eta_h(M_h(\lambda),\Omega)$ and the error $D(M(\lambda), M_h(\lambda))$ over mesh $\mathcal{T}_h$ as follows:

$$\eta_h(M_h(\lambda),\Omega) = \sum_{l=1}^{q} \eta_h(u_{h,l},\Omega), \tag{6.1}$$

$$D(M(\lambda), M_h(\lambda)) = \sum_{l=1}^{q} \|u_l - E_h u_l\|_{a,\Omega}. \tag{6.2}$$

We see from the proof of Lemma 3.1 that

$$\eta_h(M_h(\lambda),\Omega) \approx \sum_{l=1}^{q} \eta_h(E_h u_l,\Omega).$$

Thus we get from Theorem 3.2 that

$$\eta_h(M_h(\lambda),\Omega) \approx D(M(\lambda), M_h(\lambda)).$$

Example 1 Consider the following harmonic oscillator equation, which is a simple model in quantum mechanics [18]:

$$-\frac{1}{2}\Delta u+\frac{1}{2}|x|^2u=\lambda u \ \text{ in } \mathbb{R}^3, \tag{6.3}$$

where $|x|=\sqrt{|x_1|^2+|x_2|^2+|x_3|^2}$. The eigenvalues of (6.3) are $\lambda_n=n+\frac{1}{2}$ with multiplicity $\frac{n(n+1)}{2}(n=1,2,\cdots)$, and its associated eigenfunction is $u_n=\gamma e^{-|x|^2/2}H_n(x)$ with any nonzero constant γ and $H_n(x)=(-1)^ne^{x^2}\frac{d^n}{dx^n}(e^{-x^2})$.

Since the solution of (6.3) exponentially decays, we may solve it over some bounded domain Ω. In the computation, we solve the following eigenvalue problem: find $(\lambda,u)\in\mathbb{R}\times H_0^1(\Omega)$ such that $\int_\Omega |u|^2dx=1$ and

$$\begin{cases} -\frac{1}{2}\Delta u+\frac{1}{2}|x|^2u &= \lambda u \ \text{ in } \Omega,\\ u &= 0 \ \text{ on } \partial\Omega,\end{cases}$$

where $\Omega=(-5.5,5.5)^3$. We calculate the approximation of the first two smallest eigenvalues λ_1 and λ_2 with multiplicity 1 and 3, respectively, and their corresponding eigenfunction spaces $M(\lambda_1)$ and $M(\lambda_2)$ with dimension 1 and 3, respectively.

Some cross-sections of the adaptively refined mesh constructed by the **Marking Strategy E** are displayed in Fig. 6.1, from which we observe that the mesh is denser in the center of the domain where the solutions oscillate quickly than in the domain far away from the center where the solution is smoother. This shows that our adaptively refined mesh can catch the oscillation of the solution efficiently and the a posteriori error estimators we designed are efficient. Our numerical results are presented in Fig. 6.2 and Fig. 6.3. Since the multiplicity of the first two smallest eigenvalues are 1 and 3, respectively, for the discrete problem, we calculate the first 4 eigenpairs. We see from the left figure of Figure 6.2 that the convergence curves of error for all eigenvalues by using linear finite elements are parallel to the line with slope $-\frac{2}{3}$. Besides, we also observe that the convergence curves for the second, the third and the forth eigenvalues overlap together, this coincide with the fact that the multiplicity of the second eigenvalue is 3. Meanwhile, from the left figure of Fig. 6.3 we see that by using linear finite elements, the convergence curves of the a posteriori error estimators for eigenfunction space $\eta_h(M_h(\lambda_1))$ and $\eta_h(M_h(\lambda_1))$ are parallel to the line with slope $-\frac{1}{3}$. Since $\eta_h(M_h(\lambda),\Omega)\approx D(M(\lambda),M_h(\lambda))$, we obtain that the convergence curves of error for the eigenfunction space $D(M(\lambda_1),M_h(\lambda_1))$ and $D(M(\lambda_2),M_h(\lambda_2))$ are also parallel to the line with slope $-\frac{1}{3}$. This means that the approximation of eigenvalues as well as the eigenfunction space have reached the optimal convergence rate, which coincides with our theory in Section 5. We have the similar conclusion for the quadratic finite elements from the right figures of Fig. 6.2 and Fig. 6.3.

Example 2 Consider the Schrödinger equation for hydrogen atoms:

$$\left(-\frac{1}{2}\Delta-\frac{1}{|x|}\right)u=\lambda u \ \text{ in } \mathbb{R}^3 \tag{6.4}$$

with $\int_{\mathbb{R}^3}|u|^2dx=1$. The eigenvalues of (6.4) are $\lambda_n=-\frac{1}{2n^2}(n=1,2,\cdots)$ and the multiplicity of λ_n is n^2 (see, e.g., [19]).

Since the eigenfunctions of (6.4) decay exponentially, instead of (6.4), we may solve the following eigenvalue problem: find $(\lambda,u)\in\mathbb{R}\times H_0^1(\Omega)$ such that $\int_\Omega|u|^2dx=1$ and

$$\begin{cases} \left(-\frac{1}{2}\Delta-\frac{1}{|x|}\right)u &= \lambda u \ \text{ in } \Omega,\\ u &= 0 \ \text{ on } \partial\Omega,\end{cases} \tag{6.5}$$



(a) Linear finite elements (b) Quadratic finite elements

Figure 6.1: The cross-sections of an adaptive mesh of **Example 1**



(a) Linear finite elements (b) Quadratic finite elements

Figure 6.2: The convergence curve of relative error for the eigenvalues of **Example 1**

where Ω is some bounded domain in $\mathbb{R}^3$. In our computation, we choose $\Omega = (-20.0, 20.0)^3$ and find the first 2 smallest eigenvalue approximations and their associated eigenfunction space approximations. Since the multiplicity of the n-th smallest eigenvalue is n^2, for the discrete problem of (6.5), we calculate the first 5 smallest eigenvalues and their associated eigenfunctions.

Fig. 6.4 is the cross-sections of the adaptively refined mesh constructed by **Marking Strategy E**. Similarly, we see that for both the linear finite elements and quadratic finite elements, the mesh is much denser in the center of the domain where the solution oscillates quickly than in the domain far away from the center where the solution is smooth. This means that the a posteriori error estimators we used are efficient.

The numerical results are presented in Fig. 6.5 and Fig. 6.6. Similar to **Example 1**, Fig. 6.5 shows that the convergence curve for all the eigenvalues obtained by linear finite elements and quadratic finite elements are parallel to the line with slope $-\frac{2}{3}$ and $-\frac{4}{3}$, respectively, which means all the eigenvalue approximations reach the optimal convergence rate for both linear finite element and quadratic finite element. Meanwhile, we see from Fig. 6.6 that convergence curve for the a posteriori error estimators for eigenfunction space $\eta_h(M_h(\lambda_1))$ and $\eta_h(M_h(\lambda_2))$ obtained by linear finite element are parallel to the line with slope $-\frac{1}{3}$, and those obtained by quadratic finite element are parallel to the line with slope $-\frac{2}{3}$. We observe that the approximation of eigenfunction space



(a) Linear finite elements (b) Quadratic finite elements

Figure 6.3: The convergence curves of $\eta_h(M_h(\lambda_1), \Omega)$ and $\eta_h(M_h(\lambda_2), \Omega)$ for **Example 1**



(a) Linear finite elements (b) Quadratic finite elements

Figure 6.4: The cross-section of an adaptive mesh of **Example 2**

has also reached optimal convergent rate. These results validate our theoretical results.

Example 3 Consider the following eigenvalue problem which is defined in a non-convex domain: find $(\lambda, u) \in \mathbb{R} \times H_0^1(\Omega)$ such that $\displaystyle\int_\Omega |u|^2 dx = 1$ and

$$\left\{ \begin{array}{rcl} -\dfrac{1}{2}\Delta u & = & \lambda u \ \text{ in } \Omega, \\ u & = & 0 \ \text{ on } \partial\Omega, \end{array} \right. \tag{6.6}$$

where $\Omega = (-5.0, 5.0)^3 \setminus (0, 5.0)^3$, see the left figure of Fig. 6.7 below. We observe from the numerical calculation that $\lambda_1 = 0.210651$ with multiplicity 1 and $\lambda_2 = 0.331779$ with multiplicity 2.

The surface of the adaptively refined meshes constructed by **Marking Strategy E** is shown in Fig. 6.7. Besides, some cross-sections are displayed in Fig. 6.7. We see from these figures that for both linear finite elements and quadratic finite elements, the mesh is much denser along the lines where the solution is singular than in the domain far away from the singular lines. It indicates that our error estimator and marking strategy are efficient.

Our numerical results are listed in Fig. 6.8 and Fig. 6.9. Similar to **Example 1** and **Example 2**, we can also see that the approximations of eigenvalue as well as eigenfunction have reached optimal convergence rate, which coincides with our theory in Section 5.



(a) Linear finite elements (b) Quadratic finite elements

Figure 6.5: The convergence curves of relative error for eigenvalues of **Example 2**



(a) Linear finite elements (b) Quadratic finite elements

Figure 6.6: The convergence curves of $\eta_h(M_h(\lambda_1), \Omega)$ and $\eta_h(M_h(\lambda_2), \Omega)$ for **Example 2**

7 Concluding remarks

We have studied the convergence and complexity for elliptic multiple eigenvalue problems. Now we turn to address how to apply the same arguments to the Steklov problem that consists in finding $\lambda \in \mathbb{R}$ and $u \neq 0$ such that

$$\begin{cases} -\nabla \cdot (A\nabla u) + cu & = 0 \quad \text{in} \quad \Omega, \\ (A\nabla u) \cdot \overrightarrow{n} & = \lambda uv \quad \text{on} \quad \partial\Omega, \end{cases}$$

where $\overrightarrow{n}$ is the outward unit normal vector of Ω on $\partial\Omega$.

We set $H = H^1(\Omega)$, $V_h = S^h(\Omega)$, $W = L^2(\partial\Omega)$, $b(\cdot,\cdot) = (\cdot,\cdot)_{\partial\Omega}$ and consider the non-homogeneous Neumann problem as a model problem as follows:

$$\begin{cases} Lu_i & = 0 \text{ in } \quad \Omega, \quad i = 1, \cdots, N, \\ (A\nabla u_i) \cdot \overrightarrow{n} & = f_i \text{ on } \partial\Omega. \end{cases} \tag{7.1}$$



(a) nonconvex domain Ω (b) Linear finite elements (c) Quadratic finite elements

Figure 6.7: The surface of an adaptive mesh of **Example 3**



(a) Linear finite elements (b) Quadratic finite elements

Figure 6.8: The convergence curve of relative error for eigenvalues of **Example 3**

Define the element residual $\tilde{\mathcal{R}}_T(u_{i,h})$ and the jump residual $\tilde{J}_E(u_{i,h})$ for (7.1) as follows:

$$\begin{aligned} \tilde{\mathcal{R}}_T(u_{i,h}) &= \nabla\cdot(A\nabla u_{i,h}) - cu_{i,h} \quad \text{in } T\in\mathcal{T}_h, \\ \tilde{J}_E(u_{i,h}) &= \begin{cases} [[A\nabla u_{i,h}]]_E\cdot\nu_E & \text{on } E\in\mathcal{E}_h \\ A\nabla u_{i,h}\cdot\overrightarrow{n} - f_i & \text{on } E\in\mathcal{S}_h, \end{cases} \end{aligned}$$

where $\mathcal{S}_h$ denote the set of boundary faces. For $T\in\mathcal{T}_h$, we denote the local error indicator $\tilde{\eta}_h(u_{i,h},T)$ by

$$\tilde{\eta}_h^2(u_{i,h},T) = h_T^2\|\tilde{\mathcal{R}}_T(u_{i,h})\|_{0,T}^2 + \sum_{E\in\mathcal{E}_h\cup\mathcal{S}_h, E\subset\partial T} h_E\|\tilde{J}_E(u_{i,h})\|_{0,E}^2$$

and the oscillation $\widetilde{osc}_h(u_{i,h},T)$ by

$$\widetilde{osc}_h^2(u_{i,h},T) = h_T^2\|\tilde{\mathcal{R}}_T(u_{i,h}) - \overline{\tilde{\mathcal{R}}_T(u_{i,h})}\|_{0,T}^2 + \sum_{E\in\mathcal{E}_h\cup\mathcal{S}_h, E\subset\partial T} h_E\|\tilde{J}_E(u_{i,h}) - \overline{\tilde{J}_E(u_{i,h})}\|_{0,E}^2. \quad (7.2)$$

We see that Lemma 2.5 is also valid for (7.2).



(a) Linear finite elements

(b) Quadratic finite elements

Figure 6.9: The convergence curves of $\eta_h(M_h(\lambda_1),\Omega)$ and $\eta_h(M_h(\lambda_2),\Omega)$ for **Example 3**

In context of Steklov eigenvalue problems, we define

$$\begin{aligned} \mathcal{R}_T(E_hu) &= \nabla\cdot(A\nabla E_hu)-cE_hu \quad \text{in } T\in\mathcal{T}_h,\\ J_E(E_hu) &= \begin{cases} [[A\nabla E_hu]]_E\cdot\nu_E & \text{on } E\in\mathcal{E}_h\\ A\nabla E_hu\cdot\overrightarrow{n}-\lambda^hE_hu & \text{on } E\in\mathcal{S}_h,\end{cases}\end{aligned}$$

where $\overrightarrow{n}$ denotes the outward unit normal vector on $E\in\mathcal{S}_h$. For $T\in\mathcal{T}_h$, we define the local error indicator $\eta_h(E_hu,T)$ by

$$\eta_h^2(E_hu,T)=h_T^2\|\mathcal{R}_T(E_hu)\|_{0,T}^2+\sum_{E\in\mathcal{E}_h\cup\mathcal{S}_h,E\subset\partial T}\|J_E(E_hu)\|_{0,E}^2$$

and the oscillation $osc_h(E_hu,T)$ by

$$osc_h^2(E_hu,T)=h_T^2\|\mathcal{R}_T(E_hu)-\overline{\mathcal{R}_T(E_hu)}\|_{0,T}^2+\sum_{E\in\mathcal{E}_h\cup\mathcal{S}_h,E\subset\partial T}h_E\|J_E(E_hu)-\overline{J_E(E_hu)}\|_{0,E}^2.$$

We obtain by using the same argument that Theorem 4.1 and Theorem 5.1 are valid for the Steklov problem with multiple eigenvalues.

In our numerical analysis above, for convenience, we assume that the algebraic eigenvalue problem is exactly solved and the numerical integration is exact. Indeed, the same conclusion can be expected if all the numerical errors are taken into account, including both the error resulting from the inexact solving of the algebraic eigenvalue problem and the error coming from the inexact numerical integration. Suppose (λ,u) is an eigenpair with the multiplicity of λ being q, the exact solution on mesh $\mathcal{T}_k$ are $\{(\lambda_{h,i},u_{h,i})\}_{i=1}^q$, and the the solution considering the numerical error are $\{(\hat{\lambda}_{h,i},\hat{u}_{h,i})\}_{i=1}^q$. If the numerical errors resulting from the solution of algebraic system and the numerical integration are small enough, say, satisfy

$$\sum_{i=1}^q\left(\|u_{h,i}-\hat{u}_{h,i}\|_a^2+|\lambda_{h,i}-\hat{\lambda}_{h,i}|\right)\lesssim r(h_0)\sum_{i=1}^q\eta_h^2(\hat{u}_{h,i},\Omega)$$

with $r(h_0)\ll 1$ for $h_0\ll 1$, then we have from the following triangle inequality

$$\|u_i-\hat{u}_{h,i}\|_a\le\|u_i-u_{h,i}\|_a+\|u_{h,i}-\hat{u}_{h,i}\|_a,$$

that our main results obtained in this paper hold true for inexact algebraic solution and inexact numerical integration, too.

References

[1] R.A. Adams, *Sobolev Spaces*, Academic Press, New York, 1975.

[2] I. Babuska and J.E. Osborn, *Finite element-Galerkin approximation of the eigenvalues and eigenvectors of selfadjoint problems*, Math. Comput., **52** (1989), pp. 275-297.

[3] I. Babuska and J.E. Osborn, *Eigenvalue problems, in: Handbook of Numerical Analysis, Vol. II, Finite Element Methods (Part 1) (P.G. Ciarlet and J.L. Lions, eds.)*, Elsevier, pp. 641-792.

[4] I. Babuska and M. Vogelius, *Feedback and adaptive finite element solution of one-dimensional boundary value problems*, Numer. Math., **44** (1984), pp. 75-102.

[5] R. Becker and R. Rannacher, *An optimal control approach to a posteriori error estimation in finite element methods*, Acta Numerica, **10** (2001), pp. 1-102.

[6] P. Binev, W. Dahmen, and R. DeVore, *Adaptive finite element methods with convergence rates*, Numer. Math., **97** (2004), pp. 219-268.

[7] J.M. Cascon, C. Kreuzer, R.H. Nochetto, and K.G. Siebert, *Quasi-optimal convergence rate for an adaptive finite element method*, SIAM J. Numer. Anal., **46** (2008), pp. 2524-2550.

[8] F. Chatelin, *Spectral Approximations of Linear Operators*, Academic Press, New York, 1983.

[9] P.G. Ciarlet, and J.L. Lions, *Handbook of Numerical Analysis, Vol.II, Finite Element Methods (Part I)*, North-Holland, 1991.

[10] W. Dahmen, T. Rohwedder, R. Schneider, and A. Zeiser, *Adaptive eigenvalue computation: Complexity estimates*, Numer. Math., **110** (2008), pp. 277-312.

[11] X. Dai, X. Gong, Z. Yang, D. Zhang, and A. Zhou, *Finite volume discretizations for eigenvalue problems with applications to electronic structure calculations*, Multiscale Model. Simul., **9** (2011), pp. 208-240.

[12] X. Dai, J. Xu, and A. Zhou, *Convergence and optimal complexity of adaptive finite element eigenvalue computations*, Numer. Math., **110** (2008), pp. 313-355.

[13] W. Dörfler, *A convergent adaptive algorithm for Poisson's equation*, SIAM J. Numer. Anal., **33** (1996), pp. 1106-1124.

[14] R. Durán, C. Padra, and R. Rodríguez, *A posteriori error estimates for the finite element approximation of eigenvalue problems*, Math. Mod. Meth. Appl. Sci., **13** (2003), pp. 1219-1229.

[15] E.M. Garau and P. Morin, *Convergence and quasi-optimality of adaptive FEM for Steklov eigenvalue problems*, IMA J. Numer. Anal., **31** (2011), pp. 914-946.

[16] E.M. Garau, P. Morin, and C. Zuppa, *Convergence of adaptive finite element methods for eigenvalue problems*, M^{3}AS, **19** (2009), pp. 721-747.

[17] S. Giani and I. G. Graham, *A convergent adaptive method for elliptic eigenvalue problems*, SIAM J. Numer. Anal., **47** (2009), pp. 1067-1091.

[18] W. Greiner, Quantum Mechanics. An Introduction, Springer-Verlag, Berlin, 1989.

[19] W. Greiner, Quantum Mechanics: An Introduction, 3rd ed., Springer-Verlag, Berlin, Heidelberg, 1994.

[20] L. He and A. Zhou, *Convergence and complexity of adaptive finite element methods for elliptic partial differential equations*, Inter. J. Numer. Anal. Model., **8** (2011), pp. 615-640.

[21] V. Heuveline and R. Rannacher, *A posteriori error control for finite element approximations of elliptic eigenvalue problems*, Adv. Comput. Math., **15** (2001), pp. 107-138.

[22] M.G. Larson, *A posteriori and a priori error analysis for finite element approximations of self-adjoint elliptic eigenvalue problems*, SIAM J. Numer. Anal., **38** (2000), pp. 608-625.

[23] C. Le Bris, ed., *Handbook of Numerical Analysis, Vol. X. Special issue: Computational Chemistry*, North-Holland, Amsterdam, 2003.

[24] D. Mao, L. Shen, and A. Zhou, *Adaptive finite element algorithms for eigenvalue problems based on local averaging type a posteriori error estimates*, Adv. Comput. Math., **25** (2006), pp. 135-160.

[25] R.M. Martin, *Electronic Structure: Basic Theory and Practical Method*, Cambridge University Press, Cambridge, 2004.

[26] V. Mehrmann and A. Miedlar, *Adaptive computation of smallest eigenvalues of self-adjoint elliptic partial differential equations*, Numer. Linear Algebra Appl., **18** (2011), pp. 387-409.

[27] K. Mekchay and R. Nochetto, *Convergence of adaptive finite element methods for general second order linear elliptic PDEs*, SIAM J. Numer. Anal., **43** (2005), pp. 1803-1827.

[28] P. Morin, R.H. Nochetto, and K. Siebert, *Convergence of adaptive finite element methods*, SIAM Review, **44** (2002), pp. 631-658.

[29] Y. Saad, J.R. Chelikowsky, and S.M. Shontz, *Numerical methods for electronic structure calculations of materials*, SIAM Review, **52** (2010), pp. 3-54.

[30] L. Shen and A. Zhou, *A defect correction scheme for finite element eigenvalues with applications to quantum chemistry*, SIAM J. Sci. Comput., **28** (2006), pp. 321-338.

[31] R. Stevenson, *Optimality of a standard adaptive finite element method*, Found. Comput. Math., **7** (2007), pp. 245-269.

[32] R. Stevenson, *The completion of locally refined simplicial partitions created by bisection*, Math. Comput., **77** (2008), pp. 227-241.

[33] R. Verfürth, *A Riview of a Posteriori Error Estimates and Adaptive Mesh-Refinement Techniques*, Wiley-Teubner, New York, 1996.

[34] J. Xu and A. Zhou, *Local and parallel finite element algorithms based on two-grid discretizations*, Math. Comput., **69** (2000), pp. 881-909.

[35] J. Xu and A. Zhou, *A two-grid discretization scheme for eigenvalue problems*, Math. Comput., **70** (2001), pp. 17-25.